arXiv:1204.3980v1 [math.PR] 18 Apr 2012

GENERALIZED BOOTSTRAP PERCOLATION

BÉLA BOLLOBÁS, PAUL SMITH, AND ANDREW UZZELL

Abstract. In this paper we introduce generalized bootstrap processes on $\mathbb{Z}^2$ with local, homogeneous, monotone update rules of any kind, and prove the first results about the phase transitions of these processes. Let $\mathcal{U} = \{X_1, \dots, X_m\}$ be a set of finite, non-empty subsets of $\mathbb{Z}^2 \setminus \{0\}$. Under the generalized bootstrap process for $\mathcal{U}$, there is an initial set $A_0 \subset \mathbb{Z}^2$ of 'infected' sites, and at time $t \geqslant 1$ the infected set is $A_t = A_{t-1} \cup \{x \in \mathbb{Z}^2 : X_i + x \subset A_{t-1} \text{ for some } i \in [m]\}$. Let $p_c(\mathcal{U}, t) = \inf\{p : \mathbb{P}_p(0 \in A_t) \geqslant 1/2\}$, where A_0 is a random subset of $\mathbb{Z}^2$ in which sites are infected independently with probability p. We prove that, under certain weak conditions, there exist constants $\lambda_1, \lambda_2 > 0$ and integers $\alpha_1, \alpha_2 \geqslant 1$ such that

$$\left(\frac{\lambda_1}{\log t}\right)^{1/\alpha_1} \leqslant p_c(\mathcal{U}, t) \leqslant \left(\frac{\lambda_2}{\log t}\right)^{1/\alpha_2}.$$

These are the first results of any kind on bootstrap percolation considered in this level of generality.

1. Introduction

Cellular automata are dynamical systems on graphs in which vertices have one of a finite number of states at discrete times t. The states of the vertices change according to an update rule, which is homogeneous (the same rule applies to every vertex) and local (the rule depends on only a finite neighbourhood of the vertex under consideration). These systems were first introduced by von Neumann (see [20], for example) at the suggestion of Ulam [18]. The subject of this paper is *bootstrap percolation*, a particular type of cellular automaton proposed by Chalupa, Leith and Reich in 1979 [6], who were motivated by applications to interacting particle systems in physics.

In the simplest model of bootstrap percolation, called *r-neighbour bootstrap percolation*, a subset A of the vertices of a graph G are initially *infected*, and at each time t, infected vertices stay infected, and among the uninfected vertices of G, those which have at least r infected neighbours become infected. In keeping with the percolation literature, we shall use the words 'vertex' and 'site' interchangeably. Most mathematical results in the field of bootstrap percolation have concerned the r-neighbour model, either on $\mathbb{Z}^d$ or the finite grid $[n]^d$ (see [19, 1, 17, 16, 14, 3, 2] for a selection of these results). Gravner and Griffeath [11, 12, 13] were the first to generalize the ideas from the r-neighbour process to other bootstrap processes. They studied bootstrap processes, again on integer lattices, but this time with much more general update rules, called *threshold dynamics*. These are defined by a neighbourhood $\mathcal{N}$, which is a finite subset of $\mathbb{Z}^d \setminus \{0\}$, and a positive integer threshold r. Given an initial set $A := A_0 \subset \mathbb{Z}^d$, for $t \geqslant 0$ we set

$$A_{t+1} := A_t \cup \{x \in \mathbb{Z}^d : |\mathcal{N} + x| \geqslant r\}.$$

The research of the first author was supported in part by NSF grants CNS-0721983, CCF-0728928 and DMS-0906634, and ARO grant W911NF-06-1-0076. The research of the third author was supported in part by NSF grant DMS-0906634 and ARO grant W911NF-06-1-0076. The authors are grateful to Yuval Peres and other members of the Theory Group at Microsoft Research, Redmond, where much of the research in this paper was carried out.

Thus, a site x becomes infected when at least r sites in the neighbourhood $\mathcal{N} + x$ are infected. When $\mathcal{N} = \{\pm e_1, \ldots, \pm e_d\}$, where the e_i are the standard basis vectors in $\mathbb{R}^d$, this is precisely the r-neighbour model. The threshold dynamics models studied by Gravner and Griffeath, and later Duminil-Copin and Holroyd [7], had a number of additional constraints. The first of these constraints was that the process should be critical, meaning roughly that growth is neither too easy (there exist finite sets with infinite closure) nor too hard (there exist closed cofinite sets). This is a natural assumption: if growth is not critical, then the behaviour of the critical probability is genuinely different, as we shall see later in the paper. However, the other constraints, which were that the process should be symmetric ($x \in \mathcal{N}$ implies $-x \in \mathcal{N}$) and balanced (meaning roughly that the two hardest directions in which to grow are equally hard), are much less natural.

Our aim in this paper is to go considerably further: rather than one rather special model, we shall take all local, homogeneous, monotone models. This means that we shall fix an s and take all neighbourhoods of x which consist only of sites at distance at most s from x; for each of these we say whether or not x becomes infected given that all the sites in the neighbourhood are infected, and we do this such that the neighbourhoods which infect x form an up-set. The same rule is then applied to every site x.

Formally, let $m \in \mathbb{N}$, and for each $i \in [m]$, let X_i be a finite, non-empty subset of $\mathbb{Z}^2 \setminus \{0\}$. Let $\mathcal{U} = \{X_1, \ldots, X_m\}$. We call $\mathcal{U}$ an *update family* and each X_i an *update rule.* Let $\Lambda \subset \mathbb{Z}^2$ and let $A \subset \Lambda$. In bootstrap percolation on Λ with update family $\mathcal{U}$ and initial set $A = A_0$, we set

$$A_{t+1} := A_t \cup \{x \in \Lambda : \text{there exist } i \in [m] \text{ such that } X_i + x \subset A_t\}.$$

Roughly speaking, there is a set $A \subset \Lambda$ of initially infected sites and at each time step, whenever a translate of a configuration X_i by an element of $\mathbb{Z}^2$ is completely infected, the corresponding translate of the origin also becomes infected. We call this a *generalized bootstrap process.* (The indefinite article indicates the dependence on $\mathcal{U}$; since $\mathcal{U}$ will always be understood, from now on we shall write *the generalized bootstrap process.*) The *closure of* A *under* $\mathcal{U}$ is defined to be $[A] := \cup_{t=0}^{\infty} A_t$. When the closure of A is the whole of Λ, we say that A *percolates* Λ *under* $\mathcal{U}$, or more usually, when Λ and $\mathcal{U}$ are understood from the context, we say simply that A *percolates.*



FIGURE 1. An example of an update family $\mathcal{U}$ consisting of six rules. Under the generalized bootstrap process with this update family, if there is a translate of one of the rules such that all of the black dots are infected, then the white dot will become infected too.

We are primarily interested in the case when the initial configuration A is chosen at random. Fix $p \in [0,1]$ and let Π_p be a random subset of $\mathbb{Z}^2$ in which each site is included independently with probability p. Define the initial set A to be $\Pi_p \cap \Lambda$. Let $P_p(\mathcal{U}, \Lambda)$ be the probability that A percolates Λ under $\mathcal{U}$ when A is chosen with this distribution. Closely related to $P_p(\mathcal{U},\Lambda)$, and the parameter that we shall mostly be concerned with, is the probability $P_p(\mathcal{U},t)$ that the origin is infected by time t under the generalized bootstrap process with initial set $A = \Pi_p$. The *critical probability for infecting the origin by time* t is

$$p_c(\mathcal{U},t) = \inf\{p : P_p(\mathcal{U},t) \geqslant 1/2\}.$$

As we shall show, the behaviour of bootstrap percolation under $\mathcal{U}$ is largely governed by the directions of certain unit vectors in the plane, which are normal to the boundaries of certain half-planes. Before we make this statement precise, let us set out some standard notation. Throughout, $\|\cdot\|$ will mean $\|\cdot\|_2$, the Euclidean distance, and $\langle\cdot,\cdot\rangle$ will mean the Euclidean inner product. Let $S^1 = \{x \in \mathbb{R}^2 : \|x\| = 1\}$ be the unit circle in $\mathbb{R}^2$. The definitions we give next are the most important definitions of the paper.

Let $u \in S^1$ and let $a \in \mathbb{R}^2$. We define $\mathbb{H}_{u,a}$ to be the discrete half-plane $\{z \in \mathbb{Z}^2 : \langle z - a, u\rangle < 0\}$, and we set $\mathbb{H}_u = \mathbb{H}_{u,0}$. A unit vector $u \in S^1$ is a *stable direction for* $\mathcal{U}$ if $[\mathbb{H}_u] = \mathbb{H}_u$. We denote by $\mathcal{S}$ the set of all stable directions for $\mathcal{U}$ and we call $\mathcal{S}$ the *stable set for* $\mathcal{U}$. If $u \in \mathcal{S}^c = S^1 \setminus \mathcal{S}$, then we call u an *unstable direction for* $\mathcal{U}$.

Under two-neighbour bootstrap percolation on $\mathbb{Z}^2$, the stable directions are $(1,0)$, $(0,1)$, $(-1,0)$ and $(0,-1)$. Closed sets are unions of (solid) rectangles mutually at ℓ_1 distance at least 2 whose sides are orthogonal to stable directions. One of the difficulties we encounter is that we do not have an analogous useable description of closed sets under the generalized bootstrap process. However, we shall need at least an approximate description. Let us call these approximations to set closures *strongly closed sets*. (We shall not use this expression beyond the next two paragraphs, but it has the advantage over what we shall call them later that it does not pre-empt the definition.) What properties would we like strongly closed sets to have? From the point of view of a lower bound for the critical probability, we would like the sets to be growth limiting; thus, if A is a subset of a strongly closed set, then the closure of A should be contained within the original set. (This means that strongly closed sets should be closed.) On the other hand, when proving an upper bound for the critical probability, we would like to be able to use the sets as foundations for growth, meaning that strongly closed sets (of any size) should be able to grow into larger strongly closed sets by the addition of a bounded number of additional sites. Of course, we are free to choose different definitions of strongly closed sets to statisfy each of these two properties, but it turns out that one definition is enough.

We now formalize these ideas. Let $\mathcal{U}$ be an update family and $\mathcal{S} = \{u_i : i \in I\}$ its set of stable directions. Let $J \subset I$ and for each $j \in J$ let $a_j \in \mathbb{R}^2$. An *ocean for* $\mathcal{S}$ is a set D of the form

$$D = \bigcap_{j \in J} \mathbb{H}_{u_j,a_j}.$$

In other words, D is an intersection of stable half-planes. When D is finite we call it a *droplet for* $\mathcal{S}$. We shall often not mention $\mathcal{S}$ when it is understood from the context. By the definition of a stable direction, we always have $[D] = D$ whenever D is an ocean.

There is a good reason why we define *oceans for* $\mathcal{S}$ and *droplets for* $\mathcal{S}$, rather than *oceans for* $\mathcal{U}$ and *droplets for* $\mathcal{U}$. We want the definition of a droplet to be such that we can use droplets as a base from which it is possible to grow — by this we mean that it should be possible to increase significantly the size of the set closure by the addition of a small number of extra infected sites. In two-neighbour bootstrap percolation, the closure of a rectangle and an additional site on one of its sides (so the ℓ_1 distance from the site to the rectangle is 1) is the smallest rectangle that contains the original rectangle and the extra site. So we could think of rectangles, with the help of a few additional sites, as growing into larger rectangles. Similarly, in the generalized setting, we want droplets to grow into droplets. However, it is not the case that droplets for $\mathcal{S}$ grow into droplets for $\mathcal{S}$; we find that, even when several additional sites are infected on one side of a droplet, the infection does not quite spread all the way along the side. As the droplet grows, the result is not another droplet, but instead a set of sites that resembles a droplet with its corners rounded off. We avoid this problem by defining a new set $\mathcal{S}_1$, which we obtain from $\mathcal{S}$ by adding a number of *quasi-stable directions.* It will turn out that, with a suitable definition of these quasi-stable directions, droplets for $\mathcal{S}_1$ *do* grow into droplets for $\mathcal{S}_1$. So we shall have reason to talk about oceans and droplets for both $\mathcal{S}$ and $\mathcal{S}_1$ within the context of the same update family $\mathcal{U}$.

Suppose there does not exist a droplet for $\mathcal{S}$. Intuitively we would expect this to be true if and only if there exist finite sets with infinite closures; this latter property suggests that the nature of $P_p(\mathcal{U}, t)$ in the case when there does not exist a droplet for $\mathcal{S}$ should be very different to the nature of $P_p(\mathcal{U}, t)$ in cases such as two-neighbour bootstrap percolation. Later we shall prove an appropriately formalized version of this statement. In fact, update families exhibit fundamentally different behaviour according to which of three different classes they belong. We say that $\mathcal{U}$ is

(i) *supercritical* if there does not exist a droplet for $\mathcal{U}$;
(ii) *critical* if there does exist a droplet for $\mathcal{U}$ and $|\mathcal{S}| < \infty$; and
(iii) *subcritical* if there does exist a droplet for $\mathcal{U}$ and $|\mathcal{S}| = \infty$.

(These definitions are similar but not identical to those of Gravner and Griffeath [11, 12, 13], and Duminil-Copin and Holroyd [7].) By far the most important of these families is the critical family, which includes the two-neighbour model, the modified two-neighbour model, threshold dynamics, the Froböse model [10], any many others.

The main theorems we prove are as follows.

Theorem 1. *Let $\mathcal{U}$ be a critical update family. Then exists a positive integer α_1 and a positive real λ_1 such that whenever $p = p(t)$ satisfies*

$$p \leqslant \left(\frac{\lambda_1}{\log t}\right)^{\frac{1}{\alpha_1}},$$

we have $P_p(\mathcal{U}, t) \to 0$ as $t \to \infty$.

Theorem 2. *Let $\mathcal{U}$ be a critical update family. Then exists a positive integer α_2 and a positive real λ_2 such that whenever $p = p(t)$ satisfies*

$$p \geqslant \left(\frac{\lambda_2}{\log t}\right)^{\frac{1}{\alpha_2}},$$

we have $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$.

Corollary 3. *Let $\mathcal{U}$ be a critical update family. Then exist positive integers α_1, α_2 and positive reals λ_1, λ_2 such that*

$$\left(\frac{\lambda_1}{\log t}\right)^{\frac{1}{\alpha_1}} \leqslant p_c(\mathcal{U}, t) \leqslant \left(\frac{\lambda_2}{\log t}\right)^{\frac{1}{\alpha_2}}. \qquad \square$$

Roughly speaking, we obtain $\alpha_1 = \alpha_2$ when it is equally difficult to grow in all stable directions. For example, in two-neighbour bootstrap percolation, no matter which side of a droplet we pick, we always need exactly one extra site to grow along the side, so in that case we obtain $\alpha_1 = \alpha_2$. We also obtain results for $P_p(\mathcal{U}, \Lambda)$ analogous to Theorems 1 and 2, but we defer their precise statements until we have defined $\mathcal{S}_1$.

Returning to the example update family $\mathcal{U}$ in Figure 1, in the proofs of Theorems 1 and 2 the values of α_1 and α_2 we obtain for this example are both equal to 1, so the theorems give

$$p_c(\mathcal{U}, t) = \Theta\left(\frac{1}{\log t}\right).$$

Thus, although generalized bootstrap percolation under $\mathcal{U}$ is far more complicated than two-neighbour bootstrap percolation, our theorems prove that the critical probability occurs at the same order of magnitude.

The rest of this paper is organized as follows. In Section 3 we prove an update family is supercritical if and only if there exists a finite subset of $\mathbb{Z}^2$ with infinite closure under $\mathcal{U}$. (One of these implications is trivial.) We also prove stronger results in some more specific cases. We then apply these results to the probabilistic setting to deduce bounds on $P_p(\mathcal{U}, t)$. The main results are proved in Section 4 and concern critical update families. Sections 4.1 and 4.2 deal with Theorem 1, the lower bound. It is a feature of almost all proofs in probabilistic bootstrap percolation that most of the proof concerns deterministic properties of the process, and so it is here. First, in Section 4.1, we prove deterministic results about a process related to the generalized bootstrap process. Then in Section 4.2 we use these results to prove Theorem 1. Sections 4.3, 4.4 and 4.5 deal with Theorem 2, the upper bound. In Section 4.3 we formalize the notion of quasi-stability from the discussion above, and in Section 4.4 we prove results about the growth of quasi-droplets. Then in Section 2 we apply these deterministic results to the probabilistic setting to prove Theorem 2. As mentioned above, we also obtain results for $P_p(\mathcal{U}, \Lambda)$ analogous to Theorems 1 and 2, which we prove in Section 4.6. Finally, in Section 5, we discuss open problems and conjectures.

2. Stable sets and stable half planes

In this section we derive a number of basic properties of the stable set and introduce some important definitions relating to the geometry of the stable set.

We define $\ell^*_{u,a}$ to be the line $\{x \in \mathbb{R}^2 : \langle x - a, u\rangle = 0\}$, $\ell_{u,a}$ to be the *lattice line* $\ell^*_{u,a} \cap \mathbb{Z}^2$, and we set $\ell^*_u = \ell^*_{u,0}$ and $\ell_u = \ell_{u,0}$. Note that if ℓ^*_u has irrational slope, then $\ell_u = \{0\}$.

Lemma 4. *If u is an unstable direction for $\mathcal{U}$, then $[\mathbb{H}_u] = \mathbb{Z}^2$.*

Proof. Let u be unstable. Then there exists $a \in A_1 \setminus A_0$ and $X \in \mathcal{U}$ such that $X + a \subset \mathbb{H}_u$. But then $X \subset \mathbb{H}_u$, since if $x \in X$, we have $\langle x + a, u\rangle < 0$, hence

$$\langle x, u\rangle \leqslant \langle a, u\rangle + \langle x, u\rangle < 0,$$

where the first inequality follows because $a \notin \mathbb{H}_u$.

So $0 \in A_1$, and by translation $\ell_u \subset A_1$. We claim that there exists a site b such that $\langle b, u\rangle > 0$ and $\mathbb{H}_{u,b} \subset A_1$. If ℓ^*_u has rational (or infinite) slope, then the set of non-empty lattice lines that are parallel to ℓ^*_u is discrete, so there is one such that is nearest to ℓ^*_u and not contained in $\mathbb{H}_u$, containing a site b, say. Then $\mathbb{H}_{u,b} = \mathbb{H}_u \cup \ell_u \subset A_1$. Otherwise, ℓ^*_u has irrational slope. In this case, let

$$\delta = \sup\{\lambda \in \mathbb{R} : X \subset \mathbb{H}_{u,-\lambda u}\}.$$

X is finite and does not intersect ℓ_u, so $\delta > 0$. Thus, if $x \in \mathbb{H}_{u,\delta u}$, then $X + x \subset \mathbb{H}_u$, so we have $\mathbb{H}_{u,\delta u} \subset A_1$. The line $\ell^*_{u,\delta u}$ may not contain any sites, but the lines parallel to ℓ^*_u that do contain sites are dense in the plane, so certainly there exists a site b with $\langle b, u\rangle > 0$ such that $\mathbb{H}_{u,b} \subset A_1$.

Since $\mathbb{H}_{u,b}$ has a site on its boundary, namely b, it is congruent to $\mathbb{H}_u$, so an inductive argument gives $\mathbb{H}_{u,tb} \subset A_t$ for all positive integers t. Thus, since $\langle b, u\rangle > 0$, we have that the closure of A contains $\bigcup_t \mathbb{H}_{u,tb} = \mathbb{Z}^2$. □

Let $\mathbb{T}$ be the torus $\mathbb{R}/2\pi\mathbb{Z}$. We shall often need to change between elements of S^1 and elements of $\mathbb{T}$, so we define the following natural bijection. Let $u : \mathbb{T} \to S^1$ be the function

$$u(\theta) = (\cos\theta, \sin\theta),$$

and let $\theta : S^1 \to \mathbb{T}$ be the inverse function, so that if $u = u(\theta)$ then $\theta(u) = \theta$.

Lemma 5. *$\mathcal{S}^c$ is a finite union of open intervals in the unit circle, and hence $\mathcal{S}$ is a closed subset of the unit circle.*

Proof. For each $i \in [m]$, let

$$T_i := \{u \in \mathcal{S}^c \,:\, X_i \subset \mathbb{H}_u\}.$$

We think of T_i as the set of directions *destabilized* by the update rule X_i. If $u \in \mathcal{S}^c$ then there must exist i such that $X_i \subset \mathbb{H}_u$, which implies that

$$\mathcal{S}^c = \bigcup T_i \tag{1}$$

Therefore it is sufficient to prove that each T_i is an open interval in $\mathbb{T}$.

First we prove that T_i is open. Let $u \in T_i$. Then $X_i \subset \mathbb{H}_u$, or in other words, $\langle x, u\rangle < 0$ for all $x \in X_i$. So there exists $\delta > 0$ such that $\langle x, u\rangle < -\delta$ for all $x \in X_i$. Let $M = \max\{\|x\| \,:\, x \in X_i\}$ and let $u' \in S^1$ be such that $\|u' - u\| < \delta/M$. Then, for $x \in X_i$,

$$\langle u', x\rangle = \langle u, x\rangle + \langle u' - u, x\rangle \leqslant \langle u, x\rangle + \|u' - u\|\|x\|$$

by the Cauchy-Schwarz inequality. Therefore, we have

$$\langle u', x\rangle < -\delta + \delta = 0,$$

so $u' \in D_i$.

Next we prove that T_i is an interval. As noted above,

$$T_i = \bigcap_{x \in X_i} \{u : \langle x, u\rangle < 0\}.$$

That is, T_i is an intersection of intervals of length less than π, so T_i is an interval. □

Lemma 5 justifies the following definitions. Given an update rule X, let

$$\Theta(X) = \{\theta \,:\, X \subset \mathbb{H}_{u(\theta)}\},$$

and let

$$T(X) = \{u \in S^1 \,:\, X \subset \mathbb{H}_u\}.$$

Also, define $\theta^-(X) = \inf \Theta(X)$ and $\theta^+(X) = \sup \Theta(X)$. (Note that inf and sup are well-defined on the natural linear ordering on $\Theta(X)$ when considered as a subset of $\mathbb{T}$.) By Lemma 5, we can alternatively write $\Theta(X)$ as the interval $(\theta^-(X), \theta^+(X))$. When $X = X_i$, we write $\Theta_i = \Theta(X_i)$, $T_i = T(X_i)$, $\theta_i^- = \theta^-(X_i)$ and $\theta_i^+ = \theta^+(X_i)$.

Lemma 6. *Let $\mathcal{S}$ be finite and non-empty, and let $u \in \mathcal{S}$. Then ℓ_u contains at least two sites.*

Equivalently, ℓ_u contains infinitely many sites, and also equivalently, ℓ_u^* has either rational or infinite slope.

Proof. Since $\mathcal{S}$ is finite, we deduce from Lemma 5 and the definitions just given that, if u is a stable direction, then there exists i such that $u = u(\theta_i^-)$ (equivalently, there exists j such that $u = u(\theta_j^+)$). It therefore suffices to check that, for each i, the line $\ell_{u(\theta_i^-)}$ contains at least two sites. Let $\phi = \theta_i^-$, so by definition,

$$\phi = \inf \{\theta \in (\theta_i - \pi/2, \theta_i + \pi/2) \,:\, X_i \subset \mathbb{H}_{u(\theta)}\}.$$

If $\ell_{u(\phi)}$ does not contain a non-zero site, then neither does $\ell^*_{u(\phi)}$, so the latter does not intersect the closed set X_i, and therefore there exists $\delta > 0$ such that $\ell^*_{u(\phi-\epsilon)} \cap X_i = \emptyset$ for all $0 \leqslant \epsilon < \delta$, contradicting the definition of ϕ. $\square$

This lemma justifies the following definition, which generalizes the definition of *lines of direction* u in [7]. (In fact, the definition is the same, but the set of vectors to which the definition applies is more general.) Let $u \in \mathcal{S}$ and consider the lines orthogonal to u that intersect $\mathbb{Z}^2$. These lines are parallel, discrete (since ℓ_u has rational or infinite slope), and there is a natural integer-valued indexing for them. Let $\ell_u(0)$ be the line intersecting the origin, and for each $i \in \mathbb{Z}$, let $\ell_u(i)$ be the ith line in direction u. Let $\ell^*_u(i)$ be the line in $\mathbb{R}^2$ that contains $\ell_u(i)$. Analogously to these we define the ith half-plane in direction u to be $\mathbb{H}_u(i) := \mathbb{H}_{u,a}$ where a is any element of $\ell_u(i)$. We also define δ_u to be the orthogonal distance between consecutive lines of direction u and η_u to be the distance between consecutive elements of ℓ_u. We observe that $\delta_u \eta_u = 1$.

3. Supercritical families

Recall that supercritical families are those for which there does not exist a droplet. The majority of this section is concerned with proving deterministic results about supercritical families of the following nature. We show that whenever $\mathcal{U}$ is supercritical, the closure of any sufficiently large ball in $\mathbb{Z}^2$ contains an infinite subset of $\mathbb{Z}^2$. In Proposition 9 we show that if $\mathcal{S}$ is empty then the closure of any sufficiently large ball is the whole of $\mathbb{Z}^2$. In Proposition 10 we show that if $|\mathcal{S}| = 1$ then the closure of any sufficiently large ball is essentially a minimal half-plane containing that ball (a little more precisely, we show that the closure is essentially *the* minimal ocean containing that ball). In Proposition 11 we deal with general supercritical families, assuming nothing about $\mathcal{S}$, and show that the closure of any sufficiently large ball contains a one-way infinite cylindrical subset of $\mathbb{Z}^2$. None of these results are surprising: each one says that the closure of a sufficiently large set is (or contains) the maximal set that it could conceivably be (or contain).

In addition to these results, we prove that the (Euclidean) diameter of the set of infected sites increases at least linearly in time. It turns out that this property will be key to proving the one probabilistic result of this section, Theorem 13, which will say that if $\mathcal{U}$ is supercritical and $p \geqslant t^{-1/\beta}$ for some positive integer β, then $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$. This is very different in nature to the Theorem 2, which says in the critical case that if $p \geqslant (\lambda_2/\log t)^{1/\alpha_2}$ for some constants λ_2 and α_2, then $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$ (and Theorem 1 says that this is essentially the right sort of function of t for p). The result in this section hinges crucially on the existence of finite sets with infinite closure. In fact, to prove this theorem we only use Proposition 11, but we include Propositions 9 and 10 for completeness.

Given $\theta \in \mathbb{T}$, let $\mathcal{U}(\theta)$ be the set of all update rules X that destabilize $u(\theta)$, so $X \in \mathcal{U}(\theta)$ if and only if $X \subset \mathbb{H}_{u(\theta)}$. Observe that $\mathcal{U}(\theta)$ is empty precisely when $u(\theta)$ is a stable direction. Let $R_\theta(X)$ be the smallest closed rectangular subset of $\mathbb{R}^2$ with sides parallel or perpendicular to $u(\theta)$ that contains X. Let $l(\theta)$ be the maximum over $X \in \mathcal{U}(\theta)$ of the lengths of the sides of $R_\theta(X)$, and let $\epsilon(\theta)$ be the maximum over $X \in \mathcal{U}(\theta)$ of the distance from the origin to $R_\theta(X)$. There are a finite number of update rules, so there exists an l such that $l(\theta) \leqslant l$ for all θ. Since update rules are finite and $\mathbb{H}_{u(\theta)}$ does not intersect the line through the origin perpendicular to $u(\theta)$, $\epsilon(\theta)$ is strictly greater than 0 for all θ.

The following two lemmas are at the heart of Propositions 9, 10 and 11. The purpose of the lemmas is to establish that away from the stable directions, $\epsilon(\theta)$ is bounded away from 0 uniformly in θ. This tool will enable us to view the boundary of a sufficiently large ball locally like a half-plane.

Given a metric space (Y,d), let $B(x,r) = \{y \in Y : d(x,y) \leqslant r\}$ and $\tilde{B}(x,r) = \{y \in Y : d(x,y) < r\}$.

Lemma 7. *Let Y be a metric space and K a compact subspace of Y such that*

$$K \subset \bigcup_{i\in I} \tilde{B}(x_i, r_i).$$

Then there exists $\delta > 0$ such that

$$K \subset \bigcup_{i\in I} \tilde{B}(x_i, r_i - \delta). \tag{2}$$

Proof. Suppose (2) does not hold for any $\delta > 0$. Let $W_k = \cup_{i\in I}\tilde{B}(x_i, r_i - 1/k)$. Then, for each $k = 1, 2, \ldots$, there exists $\phi_k \in K \backslash W_k$. Since K is compact, there exists a subsequence $(\phi_{k_j})_{j=1}^\infty$ of $(\phi_k)_{k=1}^\infty$ which converges to some $\phi \in K$. Now $K \subset \cup_{k=1}^\infty W_k$, so there exists k_0 such that $\phi \in W_k$ for all $k \geqslant k_0$. But the W_k are open and nested, so there exists $\delta' > 0$ such that $\tilde{B}(\phi, \delta') \subset W_k$ for all $k \geqslant k_0$, contradicting the convergence of $(\phi_{k_j})_{j=1}^\infty$ to ϕ. So we have $\delta > 0$ such that (2) holds. □

Lemma 8. *Let $\hat{\mathbb{T}}$ be a closed subset of $\mathbb{T}$ such that $u(\hat{\mathbb{T}}) \cap \mathcal{S} = \emptyset$. Then there exists $\epsilon > 0$ such that $\epsilon(\theta) \geqslant \epsilon$ for all $\theta \in \hat{\mathbb{T}}$.*

Proof. The condition $u(\hat{\mathbb{T}}) \cap \mathcal{S} = \emptyset$ is equivalent to

$$\hat{\mathbb{T}} \subset \bigcup_{i=1}^m (\theta_i^-, \theta_i^+).$$

$\hat{\mathbb{T}}$ is a closed subset of $\mathbb{T}$, which is itself compact, so $\hat{\mathbb{T}}$ is compact. Therefore by Lemma 7 there exists $\delta > 0$ such that

$$\hat{\mathbb{T}} \subset \bigcup_{i=1}^m (\theta_i^- + \delta, \theta_i^+ - \delta).$$

Given $\theta \in \hat{\mathbb{T}}$, let i be such that $\theta \in (\theta_i^- + \delta, \theta_i^+ - \delta)$, and take $X(\theta) = X_i$. With these choices of update rules for the angles θ we have

$$X(\theta) \subset \mathbb{H}_{u(\theta_i^-)} \cap \mathbb{H}_{u(\theta_i^+)}$$

and $\|x\| \geqslant 1$ for all $x \in X(\theta)$, whereas we have

$$\ell^*_{u(\theta)} \subset \left(\mathbb{H}^*_{u(\theta_i^- + \delta)} \cap \mathbb{H}^*_{u(\theta_i^+ - \delta)}\right)^c.$$

So the distance from x to $\ell^*_{u(\theta)}$ is at least $\sin\delta > \delta/2$ for all $x \in X(\theta)$ (see Figure 2). Therefore we may take ϵ to be $\delta/2$. □

Throughout the remainder of this section, we take the metric space (Y,d) to be $\mathbb{Z}^2$ with the Euclidean metric. For short we write $B(r)$ to mean $B(0,r)$.

The proofs of the next three propositions are longer than the proofs of Lemmas 7 and 8, but the details are mostly technical.



FIGURE 2. $X(\theta)$ is a subset of $\mathbb{H}_{u(\theta_i^-)} \cap \mathbb{H}_{u(\theta_i^+)}$, which consists of the sites inside the shaded area. $\ell^*_{u(\theta)}$ is contained in the complement of the shaded area widened by an angle of δ at either side. The distance between these two regions is at least $\sin\delta$.

Proposition 9. *Suppose $|\mathcal{S}| = 0$. Then there exists r_0 such that $[B(r)] = \mathbb{Z}^2$ for all $r \geqslant r_0$. Furthermore, there exists $s > 0$ such that if $r \geqslant r_0$ and $A_0 = B(r)$, then $B(r + st) \subset A_t$ for all $t \geqslant 0$.*

The final statement implies that growth of the ball happens at least linearly in time. As stated in the introduction to this section, this is the key property for the probabilistic corollary.

Proof. The aim of the proof is to show that if the radius of the ball is sufficiently large, and hence its curvature sufficiently small, then it looks locally like a half-plane at points on the boundary (this is what we want: if the boundary of the ball *were* flat, then we would be done, at least if we only wanted the closure result). A little more precisely, we show that if the radius r is sufficiently large, then for all directions u, the boundary of the ball near ru looks like a half plane, so if x is a point just outside the boundary of the ball in direction u and X is the (or a) rule that destabilizes u, then $X + x$ is contained within the ball. This will give the first result. To get the linearity result we need a little refinement to show that $X + x$ is not just inside $B(r)$ but at least a constant distance from the boundary of $B(r)$.

Since $\mathcal{S}$ is empty, $u(\theta)$ is not a stable direction for any $\theta \in \mathbb{T}$, so $\mathcal{U}(\theta)$ is non-empty for all $\theta \in \mathbb{T}$. We apply Lemma 8 with $\hat{\mathbb{T}} = \mathbb{T}$ to get $\epsilon > 0$ such that $\epsilon(\theta) \geqslant \epsilon$ for all $\theta \in \mathbb{T}$. Thus, given a direction θ, there exists an update rule $X(\theta) \in \mathcal{U}(\theta)$ contained within a rectangle $R_\theta(X(\theta))$, with sides parallel and perpendicular to $u(\theta)$, side lengths at most l, and distance from the origin at least ϵ.

Recall that we want, for sufficiently large r, that if $x = \|x\|u(\theta)$ lies outside of $B(r)$, but lies sufficiently close to the boundary of $B(r)$, then $X(\theta) + x$ is contained in $B(r)$. More precisely, we require r_0 and $s > 0$ with the property that, for all $r \geqslant r_0$, if $\|x\| \leqslant r + s$ then $(R_\theta(X(\theta)) + x) \cap \mathbb{Z}^2 \subset B(r)$. Take

$$r_0 = \frac{l^2}{\epsilon} + \frac{\epsilon}{4},$$

so $r_0^2 = (r_0 - \epsilon/2)^2 + l^2$. If $r \geqslant r_0$, then $r^2 \geqslant (r - \epsilon/2)^2 + l^2$. Suppose that $\|x\| \leqslant r + \epsilon/2$ and let $y \in X(\theta)$. We must show that $\|x + y\| \leqslant r$. But this is the case, since

$$\|x + y\|^2 \leqslant (\|x\| - \epsilon)^2 + l^2 \leqslant (r - \epsilon/2)^2 + l^2 \leqslant r^2.$$

We conclude that $s = \epsilon/2$ will do, and if $A_0 = B(r)$ and $r \geqslant r_0$, then inductively $B(r+st) \subset A_t$ for all $t \geqslant 0$. ☐

Proposition 10. *Suppose* $|\mathcal{S}| = 1$ *and let* $\hat{u}$ *be the unique stable direction. Then there exists* $r_0 > 0$ *such that for all* $r \geqslant r_0$ *the following holds. Let* k *be the minimal integer such that* $r\hat{u} \in \mathbb{H}_{\hat{u}}^*(k)$. *Then*

$$\mathbb{H}_{\hat{u}}(k-1) \subset [B(r)] \subset \mathbb{H}_{\hat{u}}(k).$$

Furthermore, there exists $s > 0$ *such that if* $r \geqslant r_0$ *and* $A_0 = B(r)$, *then*

$$B(-st\hat{u}, r + st) \setminus \ell_{\hat{u}}(k-1) \subset A_t$$

for all $t \geqslant 0$.

Proof. The containment $[B(r)] \subset \mathbb{H}_{\hat{u}}(k)$ follows from the stability of $\hat{u}$. The containment $\mathbb{H}_{\hat{u}}(k-1) \subset [B(r)]$ follows from the much stronger statement in the final sentence of the Proposition, so we only have to prove that. Part of the proof follows similar lines to the proof of Proposition 9, but there is an extra difficulty here because $\epsilon(\theta)$ need not not bounded away from 0 near to $\theta(\hat{u})$.

Let $\hat{u} = u(\hat{\theta})$. Since $\hat{u}$ is the unique stable direction,

$$\bigcup_{i=1}^{m} (\theta_i^-, \theta_i^+) = \mathbb{T} \setminus \{\hat{\theta}\}.$$

Therefore there are indices a and b such that $\hat{\theta} = \theta_a^+ = \theta_b^-$. Choose arbitrarily $\theta_a \in (\theta_a^-, \theta_a^+)$ and $\theta_b \in (\theta_b^-, \theta_b^+)$, and let $\hat{\mathbb{T}} = \mathbb{T} \setminus (\theta_a, \theta_b)$. Applying Lemma 8 with this $\hat{\mathbb{T}}$ gives $\epsilon > 0$ such that $\epsilon(\theta) \geqslant \epsilon$ for all $\theta \in \hat{\mathbb{T}}$.

Fix the initial radius r_1, which should be at least r_0, a quantity to be determined later. Define the moving ball $B'(r) = B(-(r - r_1)\hat{u}, r)$, which has the property that the tangent at $r_1\hat{u}$ is fixed.

We require r_0 and $s > 0$ with the property that, for all $r \geqslant r_1 \geqslant r_0$, if $x \in B'(r+s) \setminus \ell_{\hat{u}}(k-1)$ then there exists $X \in \mathcal{U}$ such that $X + x \subset B'(r) \setminus \ell_{\hat{u}}(k-1)$. Put another way, we must show that there exist r_0 and $s > 0$ such that if x satisfies

$$\|x + (r + s - r_1)\hat{u}\| \leqslant r + s \quad \text{and} \quad x \notin \ell_{\hat{u}}(k-1) \tag{3}$$

for some $r_1 \geqslant r_0$, then there exists an update rule X such that if $y \in X$ then y satisfies

$$\|x + y + (r - r_1)\hat{u}\| \leqslant r \quad \text{and} \quad x + y \notin \ell_{\hat{u}}(k-1). \tag{4}$$

Let $v = -(r + s - r_1)\hat{u}$ and $v' = -(r - r_1)\hat{u}$. Let x have property (3) and let θ be such that $x - v = \|x - v\|u(\theta)$. We divide the remainder of the proof into three cases (albeit only two essentially different cases) according to the value of θ and show in each case that there exist r_0 and s satisfying the property for the given range of values of θ. Taking the maximum of the three bounds we obtain on r_0 and the minimum of the three bounds we obtain on s will give us the result. In the first case we assume θ is bounded away from $\hat{\theta}$ and use Lemma 8 to show that $\epsilon(\theta)$ is bounded away from zero in this region. In the second two cases, we assume θ is close to $\hat{\theta}$, and then use properties of the geometry of the rule(s) that destabilize $u(\theta)$ to prove the result.

Case 1. First suppose $\theta \in \hat{\mathbb{T}}$. We proceed as in Proposition 9, letting $X = X(\theta)$. We in fact show a stronger statement than (4), namely that

$$\|x + y + (r + s - r_1)\hat{u}\| \leqslant r - s \quad \text{and} \quad x + y \notin \ell_{\hat{u}}(k - 1) \tag{5}$$

for all $y \in X(\theta)$. This says that $x + y$ belongs to $B(v, r - s)$, while (4) says only that $x + y$ belongs to $B(v', r)$ (which contains $B(v, r - s)$). The stronger statement is more convenient to prove because $B(v, r - s)$ and $B(v, r + s)$ have the same centre (the latter being the ball than contains x).

Observe that we have $x + y \notin \ell_{\hat{u}}(k - 1)$, since $x + y \in \ell_{\hat{u}}(k - 1)$ implies

$$\cos(\hat{\theta} - \theta) \geqslant \frac{\|x + y\| - \delta_{\hat{u}}}{\|x + y\|},$$

which for $\|x\|$ sufficiently large means $\theta \notin \hat{\mathbb{T}}$.

Apply Lemma 8 with our $\hat{\mathbb{T}}$ to get $\epsilon > 0$ such that $\epsilon(\theta) \geqslant \epsilon$ for all $\theta \in \hat{\mathbb{T}}$. We choose different values for r_0 and s to those we chose in Proposition 9; this time we take

$$r_0 = \frac{3l^2}{2\epsilon} + \frac{\epsilon}{2},$$

which ensures that $(r_0 - 2\epsilon/3)^2 + l^2 = (r_0 - \epsilon/3)^2$, and we take $s = \epsilon/3$. If $r \geqslant r_1 \geqslant r_0$, then $(r - 2\epsilon/3)^2 \geqslant (r - \epsilon/3)^2 + l^2$. Therefore,

$$\|x + y - v\|^2 \leqslant (\|x - v\| - \epsilon)^2 + l^2 \leqslant (r - 2\epsilon/3)^2 + l^2 \leqslant (r - \epsilon/3)^2,$$

as required.

Case 2. Now suppose $\theta \in (\theta_a, \hat{\theta})$. We show that we can choose r_0 and s such that (4) holds when $X = X_a$. Since $x \in \mathbb{H}_{\hat{u}}(k - 1)$ and $X_a \subset \mathbb{H}_{\hat{u}}(1)$, we have $y + x \in \mathbb{H}_{\hat{u}}(k - 1)$, so $x + y \notin \ell_{\hat{u}}(k - 1)$. Therefore we just have to show that $x + y \in B'(r)$.

Let $y \in X_a$, let $\hat{u}^\perp = u(\hat{\theta} - \pi/2)$, let $x - v = x_1\hat{u} + x_2\hat{u}^\perp$, and let $x + y - v = y_1\hat{u} + y_2\hat{u}^\perp$. We are given that $x_1^2 + x_2^2 \leqslant (r + s)^2$ and must show that $(y_1 - s)^2 + y_2^2 \leqslant r$. Since $X_a \subset \mathbb{H}_{\hat{u}}(1)$, certainly $y_1 \leqslant x_1$. Furthermore, since X_a is bounded, there exists a constant c_1 such that $y_1 \geqslant x_1 - c_1$. Suppose $x_1 - 1/2 \leqslant y_1 \leqslant x_1$. We easily see that if r_1 is sufficiently large then $x_2 \geqslant 2l$, so we are free to assume that $y_2 \geqslant 0$. Then since $\|y\| \geqslant 1$, we have $0 \leqslant y_2 \leqslant x_2 - \sqrt{3}/2$, so if $s \leqslant \sqrt{3}/2$ then $x + y \in B'(r)$.

Suppose instead that $x_1 - c_1 \leqslant y_1 \leqslant x_1 - 1/2$. Since $X_a \subset \mathbb{H}_{u(\theta_a^-)}(1)$, the maximum value of $(y_1 - s)^2 + y_2$ occurs when $x + y$ lies on the line through x perpendicular to $u(\theta_a^-)$. Therefore we may assume y is of the form $\|y\|u(\theta_a^- - \pi/2)$. It follows that

$$\|x + y - v'\|^2 = \|x - v'\|^2 + 2\|x - v'\|\|y\| \cos(\theta - (\theta_a^- - \pi/2)) + \|y\|^2.$$

We have the inequalities $\theta_a^- < \theta_a \leqslant \theta$, so

$$\cos(\theta - (\theta_a^- - \pi/2)) = -\sin(\theta - \theta_a^-) \leqslant -\sin(\theta_a - \theta_a^-) \leqslant -c_2,$$

where $c_2 > 0$ is a constant. Therefore

$$\begin{aligned} \|x + y - v'\|^2 &\leqslant \|x - v'\|^2 - 2\|x - v'\|\|y\|c_2 + \|y\|^2 \\ &= (\|x - v'\| - \|y\|c_2)^2 + \|y\|^2 - (\|y\|c_2)^2 \\ &\leqslant (r + 2s - \|y\|c_2)^2 + \|y\|^2 - (\|y\|c_2)^2. \end{aligned}$$

The quantity $\|y\|$ is at least 1 and at most a constant, so taking $s = c_2/4$ we get $\|x + y - v'\|^2 \leqslant r^2$ for sufficiently large r_0.

Case 3. Finally, the case $\theta \in (\hat{\theta}, \theta_b)$ is treated similarly to Case 2.

In all three cases we have proved that (4) holds. Therefore if $A_0 = B(r_1)$ and $r_1 \geqslant r_0$, then inductively $B'(r_1 + st) \subset A_t$ for all $t \geqslant 0$. ☐

Proposition 11. *Let $\mathcal{U}$ be a supercritical update family. Then there exist constants r_0, c_1 and a unit vector $u \in \mathbb{R}^2$ with rational slope such that if $r \geqslant r_0$, then*

$$\bigcup_{\lambda \geqslant 0} B(\lambda u, r - c_1) \subset [B(r)],$$

where the union is taken over real values of λ. In particular, $|[B(r)]| = \infty$. Furthermore, there exists $s > 0$ such that if $r \geqslant r_0$ and $A_0 = B(r)$, then

$$\bigcup_{0 \leqslant \lambda \leqslant st} B(\lambda u, r - c_1) \subset A_t$$

for all $t \geqslant 0$.

Proof. First we show that there is an open interval of length at least π in $\mathbb{T}$ such that $u(\theta)$ is unstable for all θ in this interval. Equivalently, we show that there exists $\hat{\theta} \in \mathbb{T}$ such that

$$\mathcal{S} \subset \{u(\theta) : |\theta - \hat{\theta}| \geqslant \pi/2\}. \tag{6}$$

If there is no such $\hat{\theta}$, take $u(\theta_1)$, $u(\theta_2)$ and $u(\theta_3)$ stable such that $\theta_1 \in (0, \pi)$, $\theta_2 \in (\theta_1, \theta_1 + \pi)$ and $\theta_3 \in (\theta_2, \theta_2 + \pi)$. It follows that $\cap_{i=1}^3 \mathbb{H}_{u(\theta_i),u(\theta_i)}$ is a droplet, contradicting supercriticality. If there exists an open interval of values of $\hat{\theta}$ satisfying (6) then we can choose $\hat{\theta}$ such that $u(\hat{\theta})$ has rational slope. If there is no such open interval and $\hat{\theta}$ satisfies (6), then both $u(\hat{\theta} - \pi/2)$ and $u(\hat{\theta} + \pi/2)$ must be stable, so they have rational slope by Lemma 6, and therefore so does $u(\hat{\theta})$.

Let $\hat{u} = u(\hat{\theta})$, $\hat{\theta}_1 = \hat{\theta} + \pi/2$, $\hat{\theta}_2 = \hat{\theta} - \pi/2$, $\hat{u}_1 = u(\hat{\theta}_1)$ and $\hat{u}_2 = u(\hat{\theta}_2)$. We define $B''(w)$ to be a union of balls

$$B''(w) = \bigcup_{0 \leqslant \lambda \leqslant w} B(\lambda \hat{u}, r').$$

Fix $r \geqslant r_0$, where r_0 is a constant to be determined, and let r' be maximal such that r' is of the form $(k + 1/2)\delta_{\hat{u}_1}$ and $r' \leqslant r$ ($\delta_{\hat{u}_1}$ is the gap between lines of direction $\hat{u}_1$). Fix $s > 0$, a constant which is also to be determined, and let $\Delta(w) = B''(w + s) \setminus B''(w)$.

In the previous two propositions we used induction on the radius r. Here we would like to induct on λ, but at present λ is a real parameter. Therefore it would be useful to know that

$$\bigcup_{\substack{0 \leqslant \lambda \leqslant t \\ \lambda \in \mathbb{R}}} B(\lambda s \hat{u}, r') = \bigcup_{\substack{0 \leqslant \lambda \leqslant t \\ \lambda \in \mathbb{Z}}} B(\lambda s \hat{u}, r'). \tag{7}$$

when s is chosen sufficiently small. If this is the case, we would be able to assume that $\lambda \in \mathbb{Z}$ and use induction as before. Let us show that (7) holds. We must prove that $\Delta(w) \subset B((w + s)\hat{u}, r')$. Suppose this is false, and $x \in \Delta(w) \setminus B((w + s)\hat{u}, r')$ with $x = x_1 \hat{u} + x_2 \hat{u}_1$. Then

$$(r')^2 - s^2/4 < x_2^2 \leqslant (r')^2.$$

Now if

$$\frac{s^2}{4} \leqslant \frac{r_0 \delta_{\hat{u}_1}}{2} - \frac{\delta_{\hat{u}_1}^2}{16}$$

and $r' \geqslant r_0$, then we have $(r')^2 - s^2/4 \geqslant (r' - \delta_{\hat{u}_1}/4)^2$, so

$$|r' - \delta_{\hat{u}_1}/4| < |x_2| \leqslant |r'|.$$

Given that r' is of the form $(k+1/2)\delta_{\hat{u}_1}$, this means that x lies strictly between two consecutive lines of direction $\hat{u}_1$, a contradiction. So we have (7).

Therefore we must show that there exist r_0 and $s > 0$ such that for all $r \geqslant r_0$, if $x \in \Delta(w)$, then there exists $X \in \mathcal{U}$ such that $X + x \subset B''(w)$. Let $v = (w+s)\hat{u}$, $v' = w\hat{u}$, and $x - v = \|x - v\|u(\theta)$. Since every direction in $u((\hat{\theta}_2, \hat{\theta}_1))$ is unstable, there exist a and b such that $\theta_a^- < \hat{\theta}_1 \leqslant \theta_a^+$ and $\theta_b^- \leqslant \hat{\theta}_2 < \theta_b^+$. Choose arbitrarily $\theta_a \in (\theta_a^-, \hat{\theta}_1)$ and $\theta_b \in (\hat{\theta}_2, \theta_b^+)$. Let $\hat{\mathbb{T}} = [\theta_b, \theta_a]$.

We divide the remainder of the proof into three cases according to the value of θ. As in the proof of Proposition 10, the first case uses Lemma 8 to deal with values of θ in $\hat{\mathbb{T}}$, and the second two cases deal separately with with values of θ near $\hat{\theta}_1$ or $\hat{\theta}_2$.

Case 1. Suppose $\theta \in \hat{\mathbb{T}}$. Here, the proof is very similar to in Propositions 9 and 10. We start by letting $X = X(\theta)$, and we show the stronger statement that if $x \in \Delta(w)$ then $X + x \subset B((w+s)\hat{u}, r' - s)$.

Apply Lemma 8 with our $\hat{\mathbb{T}}$ to get $\epsilon > 0$ such that $\epsilon(\theta) \geqslant \epsilon$ for all $\theta \in \hat{\mathbb{T}}$. Take $s = \epsilon/2$ and

$$r_0 = \frac{l^2}{\epsilon} + \frac{3\epsilon}{4} + 1,$$

which ensures that if $r \geqslant r_0$, so $r' \geqslant r_0 - 1$, then $(r' - \epsilon/2)^2 \geqslant (r' - \epsilon)^2 + l^2$. Therefore,

$$\|x + y - v'\|^2 \leqslant (\|x - v'\| - \epsilon)^2 + l^2 \leqslant (r' - \epsilon/2)^2 = (r' - s)^2,$$

as required.

Case 2. Now suppose $\theta \in (\theta_a, \hat{\theta}_1)$. We show that $X_a + x \subset B''(w)$.

Let $y \in X_a$, let $x - v = x_1\hat{u}_1 + x_2\hat{u}$, and let $x + y - v = y_1\hat{u}_1 + y_2\hat{u}$. We are given that $x_1^2 + x_2^2 \leqslant (r')^2$. Since $X_a \subset \mathbb{H}_{\hat{u}_1}(1)$, certainly $y_1 \leqslant x_1$. Furthermore, since X_a is bounded, there exists a constant c_2 such that $y_1 \geqslant x_1 - c_2$. Suppose $x_1 - 1/2 \leqslant y_1 \leqslant x_1$. Then since $\|y\| \geqslant 1$, we have $y_2 \leqslant x_2 - \sqrt{3}/2$, so if $y_2 \geqslant 0$ and $s \leqslant \sqrt{3}/2$, then $x + y \in B''(w)$. If y_2 is negative, we still have $x + y \in B''(w)$ provided r_0 is large enough. This is because by choosing r' to be of the form $(k+1/2)\delta_{\hat{u}_1}$, we have ensured that we have as long a segment of $\ell_{\hat{u}_1}(k)$ in $B''(0)$ as we like by taking r_0 sufficiently large.

On the other hand, suppose $x_1 - c_2 \leqslant y_1 \leqslant x_1 - 1/2$. Since $X_a \subset \mathbb{H}_{u(\theta_a^-)}(1)$, the maximum value of $(y_1 + s)^2 + y_2^2$ occurs when $x + y$ lies on the line through x perpendicular to $u(\theta_a^-)$. So we may assume $y = \|y\|u(\theta_a^- - \pi/2)$. Thus

$$\|x + y - v'\|^2 = \|x - v'\|^2 + 2\|x - v'\|\|y\|\cos(\theta - (\theta_a^- - \pi/2)) + \|y\|^2.$$

We have the inequalities $\theta_a^- < \theta_a \leqslant \theta$, so $\cos(\theta - (\theta_a^- - \pi/2)) = -\sin(\theta - \theta_a^-) \leqslant -\sin(\theta_a - \theta_a^-) \leqslant -c_3$, where c_3 is a constant. Therefore

$$\begin{aligned}\|x + y - v'\|^2 &\leqslant \|x - v'\|^2 + 2\|x - v'\|\|y\|c_3 + \|y\|^2 \\ &= (\|x - v'\| - \|y\|c_3)^2 + \|y\|^2 - (\|y\|c_3)^2 \\ &\leqslant (r' + s - \|y\|c_3)^2 + \|y\|^2 - (\|y\|c_3)^2.\end{aligned}$$

Now, $\|y\|$ is at least 1 and at most a constant, so taking $s = c_3/2$ we get $\|x + y - v'\|^2 \leqslant (r')^2$ for sufficiently large r_0.

Case 3. The final case, $\theta \in (\theta_b, \hat{\theta}_2)$, is treated similarly to Case 2.

In all three cases we have proved that $x + y \in B''(w)$. Therefore if $A_0 = B''(0)$ then inductively $B''(st) \subset A_t$ for all $t \geqslant 0$. □

Corollary 12. *An update family $\mathcal{U}$ is supercritical if and only if there exists a finite set with infinite closure.* □

Theorem 13. *Let $\mathcal{U}$ be a supercritical update family. Then there exists a positive integer β such that whenever $p = p(t)$ satisfies $p \geqslant t^{-1/\beta}$ we have $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$.*

Proof. Let r_0, s and $\hat{u}$ be as in Proposition 11. If $O \notin A_t$ then by Proposition 11, A_0 does not contain a ball of radius r_0 centred at a point on the line through the origin in direction $-\hat{u}$ at distance at most st from the origin. There are at least $c_1 t$ such balls that are disjoint, where $c_1 = s/(2r_0 + 1)$. The probability that none of these balls is full is at most

$$\left(1 - p^{c_2}\right)^{c_1 t} \leqslant \exp(-c_1 t p^{c_2}),$$

where $c_2 = 4r_0^2$. If $p \geqslant t^{-1/2c_2}$ then this is at most $\exp(-c_1\sqrt{t})$. Therefore, if $\beta = 2c_2$ then

$$P_p(\mathcal{U}, t) \geqslant 1 - \exp(-c_1\sqrt{t}),$$

which tends to 1 as $t \to \infty$. □

4. CRITICAL FAMILIES

This section is the primary focus of the paper. We study critical generalized bootstrap processes with the aim of proving Theorems 1 and 2. Recall that a critical update family is an update family for which their exist droplets and such that the stable set is finite.

Theorem 1, which is the lower bound for the critical probability, is proved in Sections 4.1 and 4.2. The starting point for the proofs of both bounds is [1]. In that paper, Aizenman and Lebowitz gave the order of magnitude for the critical probability of two-neighbour bootstrap percolation on $[n]^d$. One ingredient in their proof was the observation that an internally spanned grid of side length k must contain at least k initially infected sites. They also noted that there is a certain scale (roughly $k = \log n$) at which the probability of there existing a $k \times k$ grid inside the main $n \times n$ grid with at least k initially infected sites is minimized. They are then able to show, by means of an algorithm that describes exactly how the bootstrap process runs, that if the $n \times n$ grid is internally spanned, then there must also exist a $k \times k$ grid that is internally spanned. Droplets at around the scale of $\log n$ are called *critical droplets*, because the existence of an internally spanned critical droplet is the main obstacle to percolation.

Generalized bootstrap processes present us with several problems when trying to generalize the techniques of the proof of the lower bound in [1] to the lower bound here. The most significant of these is finding an algorithm that describes exactly how the bootstrap process spreads. Instead of answering this question, we introduce a covering algorithm which dominates the bootstrap process, so that the closure of the initial set A is a subset of the final configuration under the covering algorithm. If a domain (which does not need to be a rectangle) is internally spanned under the generalized bootstrap process, then it is also covered by the algorithm. The disadvantage of this covering algorithm is that we can no longer say that there exists an internally spanned droplet at the critical scale, just that there exists a covered droplet. However, that will turn out to be enough. Key to this is a new subadditivity lemma which says, roughly speaking, that when two droplets meet, their closure cannot be too large. More specifically, it says that the projection of the closure onto any direction is at most the sum of the projections of the original droplets. In Section 4.1 we prove the subadditivity lemma and define and prove some basic properties of the covering algorithm. In Section 4.2 we put these tools together and prove Theorem 1.

The upper bound, Theorem 2, is proved in Sections 4.3, 4.4 and 4.5. Traditionally, upper bounds in bootstrap percolation are easier (in fact, often much easier) to prove than lower bounds. This does not seem to be the case here, because of the complicated geometry of the update families. For the lower bound this is less of a problem: we are able to use simple observations about stable directions to bound sets of infected sites. For the upper bound, there is no corresponding way to bound sets of infected sites *from below*; we have to be able say that certain droplets definitely are internally spanned by the bootstrap process, and for that we have to understand growth under generalized bootstrap processes. The scheme of the proof is the same as in [1]. First we find a relatively likely way for a critical droplet to be internally spanned. Then we show that we only need relatively few extra infected sites to grow from a critical droplet to the entire domain.

We encounter two main problems in the proof of the upper bound. The first, and the hardest to overcome, is the problem of growing to the corners of a droplet. When infection spreads beyond the boundary of an internally spanned droplet, what normally happens is that there is a small cluster of infected sites along one of its sides, and these infected sites (or sometimes just one site), together with the droplet itself, spread the infection along the side in both directions, so that the new internally spanned object is just the original internally spanned droplet with one of its sides extended by one unit. In other words, it is another, slightly larger, droplet. However, there are some update families under which the infection does not spread all the way to the corners of the droplet, so we get a rounding-off effect at the corners. It turns out that this can happen when some unstable directions are made unstable by more than one update rule. This might seem counter-intuitive, because we are saying that it is more difficult to grow when some directions are (in some sense) *very* unstable, but in fact this difficulty gives us a clue that this is an artificial restriction on growth. In Section 4.3 we deal with this problem by introducing *quasi-stable directions* in directions where there would otherwise be this rounding-off effect. Quasi-stable directions are not stable directions, but we treat them as such, and we grow droplets as if they were. Once we have these in place we find we are able to grow to the corners of droplets.

The second problem we encounter is that the sides of the droplets shrink. When a droplet grows along one of its side as in the previous paragraph, it may be the case that the new side has shorter length than the original side (even if we reach the corners). If it continues to grow in this way then the side we end up with may have length significantly shorter than the original side. It would then be much less likely that there still exists a cluster of sites along the side that would allow us to grow. We therefore would like a way of growing the sides of the droplets at the right relative speed so that none of the sides shrink in the long term while we are growing. In Section 4.4 we show that this is possible. We then put all of the tools for the upper bound together and prove Theorem 2 in Section 4.5.

The last section, Section 4.6, concerns versions of both results for percolation of large quasi-droplets. We have already noted in the Introduction the close relationship between the critical probabilities for hitting the origin by time n and percolation of a grid of side length n. Here we extend this relationship to generalized bootstrap processes. We obtain bounds for percolation of large quasi-droplets with the critical exponents α_1 and α_2 from Theorems 1 and 2.

In the above discussion we mentioned 'clusters of sites' several times, in the context of being able to grow along a side. In two-neighbour bootstrap percolation on $\mathbb{Z}^2$, the 'cluster' we need is just a single site. If the sites $(0,0),\dots,(k,0)$ are all infected, and a site $(a,1)$ is also infected for some $0 \leqslant a \leqslant k$, then the infection spreads to the whole of $(0,1),\dots,(k,1)$. The same is not true in the generalized setting: we may need several extra infected sites for the infection to spread along the side. (What is true though is that we only ever need a fixed number of sites, not depending on the length of the side.)

The following definition, which is one of the key definitions of the paper, is our way of formalizing this observation.

Let $\mathcal{U}$ be critical. A *breakthrough block* is a set B of the form $X \cap (\mathbb{H}_u)^c$, or a translate by an element of $\mathbb{Z}^2$ of such a set, where X is an update rule and u is a stable direction. There are no empty breakthrough blocks since if $B = X \cap (\mathbb{H}_u)^c$ is empty then X would be a subset of $\mathbb{H}_u$, so u would not be stable.



FIGURE 3. The rule X consists of the seven sites where there are large circles. The two sites where there are filled circles constitute a breakthrough block for u.

We shall also need a notion of the size of a droplet. The subadditivity lemma will involve lengths of projections of droplets, not areas, so for the lower bound it will be enough for us to know that the droplet is large in at least one direction. Let R be a (closed) rectifiable body in the plane and let $u \in S^1$. Denote by $p_E(R)$ the Euclidean perimeter of R, and by $\pi_u(R)$ the length of the orthogonal projection of R onto the line ℓ_u^*. Let $\mathrm{diam}(R)$, the *diameter* of R, be the maximal one-dimensional cross-section of R; thus,

$$\mathrm{diam}(R) := \sup_{u \in S^1} \pi_u(R).$$

A few more definitions that we shall need concern some of the geometric properties of droplets. Let

$$D = \bigcap_{i=1}^{s} \mathbb{H}_{u_i}(l_i)$$

be a droplet. We define $\mathrm{poly}(D)$ to be the closed polygonal subset of $\mathbb{R}^2$ enclosed by the lines $\ell_{u_1}^*(l_1), \dots, \ell_{u_s}^*(l_s)$, and we let $\mathrm{peri}(D) := \partial\,\mathrm{poly}(D)$, the boundary of $\mathrm{poly}(D)$. Orienting the edges of $\mathrm{peri}(D)$ anticlockwise, we define $\mathrm{side}_i(D)$ to be the $\mathbb{R}^2$ vector representing the ith side of $\mathrm{peri}(D)$. It follows trivially that

$$\sum_{i=1}^{s} \mathrm{side}_i(D) = 0.$$

Now let D be a droplet for $\mathcal{S}$. We also define p_E, π_u and diam for D; in each case, the definition is that for rectifiable bodies applied to $\mathrm{poly}(D)$. Thus, $p_E(D) := p_E(\mathrm{poly}(D))$, etc.

4.1. **Subadditivity lemmas and covered droplets.** Our aim for the first part of this section is to prove the subadditivity lemma. The lemma follows easily from a compression-type result about polygons with sides parallel to vectors from a given set, and other polygons which one could think of as a first approximation to the convex hulls of the original polygons.

Throughout this section we assume that $\mathcal{S} = \{u_1 \dots u_s\}$ is a finite subset of S^1 such that there exist droplets for $\mathcal{S}$, and that the u_i are indexed in increasing order of the angle they make with $u(0)$ anticlockwise about the origin. We do not assume $\mathcal{S}$ is the stable set of a critical update family until Lemma 17.

A (not necessarily convex) polygon P in the plane is an *$\mathcal{S}$-polygon* if for each side e_i of P, the unit vector normal to e_i oriented out of P is an element of $\mathcal{S}$. The *stable span* of R, denoted $\langle R\rangle$, is the intersection of all convex $\mathcal{S}$-polygons containing R. We define $\langle\cdot\rangle$ more generally for finite sets of points in the plane; thus, given such a set of points K, we set $\langle K\rangle = \langle \mathrm{conv}(K)\rangle \cap \mathbb{Z}^2$ (here, $\mathrm{conv}(\cdot)$ denotes the convex hull). It follows that the stable span of a droplet is just the droplet itself, and more generally, the stable span of a set of points is the minimal droplet containing that set. Note that we always have $[K] \subset \langle K\rangle$, but equality does not hold in general. (In this thesis we use *closure* and *span* to mean very different things; in most of the bootstrap percolation literature, these words are used interchangeably.)

Let P be an $\mathcal{S}$-polygon. Consider line segments ℓ with the following properties:

(i) one endpoint of ℓ is at a vertex v of P and the other is at another point u on the boundary of P;
(ii) ℓ only intersects P at u and v;
(iii) ℓ is perpendicular to some element u of $\mathcal{S}$ and the normal to ℓ in direction u is oriented towards the unbounded region of the plane.

We call such line segments *$\mathcal{S}$-rays* and we let $\mathcal{R}$ be the set of all $\mathcal{S}$-rays. Given an $\mathcal{S}$-ray ℓ, we define the *(P,ℓ)-lake*, $L(P,\ell)$, to be the finite region external to P bounded by P and ℓ. The *pre-convex hull* of P, $\mathrm{pc}(P)$, is the union of P and all (P,ℓ)-lakes; that is,

$$\mathrm{pc}(P) := P \cup \bigcup_{\ell\in\mathcal{R}} L(P,\ell).$$

It follows that $\mathrm{pc}(P)$ is an $\mathcal{S}$-polygon.

The reason for this definition is that it is easy to see that the perimeter of the pre-convex hull of an $\mathcal{S}$-polygon P is at most as long as that of the original polygon P (in fact, it is strictly shorter if there are any non-empty (P,l)-lakes), since each time we add a (P,l) lake we are replacing piecewise linear segments of the perimeter of P by a straight line segment. This is much less obviously true if one replaces 'pre-convex hull' by 'convex hull' in the previous sentence, although as we shall see, it *is* still true.

The following lemma would be better described as an observation, but it is a little technical to prove formally.

Lemma 14. *Let P be an $\mathcal{S}$-polygon. Then the sides of $\langle P\rangle$ can be obtained from those of* $\mathrm{pc}(P)$ *by translations only.*

Proof. Let u, v and w be adjacent vertices of $\mathrm{pc}(P)$. Call the vertex v *concave* if the line segment joining u to w lies externally to $\mathrm{pc}(P)$, and call it *convex* otherwise. The first step is to prove $\mathrm{pc}(P)$ does not have two consecutive concave vertices. Suppose v is concave. Let e be the line segment joining u to v and f the line segment joining v to w. Let γ be the map from edges of an $\mathcal{S}$-polygon to $[1,m]$ such that $\gamma(g) = i$ if u_i is the direction of the externally-oriented normal to the edge g. If we did not have $|\gamma(e) - \gamma(f)| = 1$, where the term inside the modulus is taken modulo s (recall that s is the number of stable directions,



FIGURE 4. The region enclosed by the solid line is an $\mathcal{S}$-polygon. The largest region enclosed by solid or dashed lines is its pre-convex hull. The largest region enclosed by solid, dashed or dotted lines is its span.

$|\mathcal{S}|$), then we would have a direction strictly between $u_{\gamma(e)}$ and $u_{\gamma(f)}$, and an $\mathcal{S}$-ray either from vertex u to side f or from vertex w to side e with the associated direction. (The $\mathcal{S}$-ray could not intersect $\mathrm{pc}(P)$ between u and f, or w and e, by the definition of the pre-convex hull.) Now suppose there are two adjacent concave vertices in $\mathrm{pc}(P)$, so we have consecutive vertices u, v, w and x with v and w concave. Let e, f and g be the edges joining u to v, v to w and w to x respectively. Then there is an $\mathcal{S}$-ray either from u to g or from x to e with direction $u_{\gamma(f)}$. So there are no two adjacent concave vertices in $\mathrm{pc}(P)$.

We apply the following algorithm to $\mathrm{pc}(P)$. If $\mathrm{pc}(P)$ does not have a concave vertex, we stop. Otherwise, let v be concave. Let u, v, w be consecutive vertices of $\mathrm{pc}(P)$ with edges e joining u to v and f joining v to w. The infinite strip enclosed by the edge e, the edge f extended beyond w, and the line parallel to f through u, is necessarily empty, otherwise there would be an additional $\mathcal{S}$-ray meeting e with outward normal direction $u_{\gamma(f)}$. For a similar reason, the infinite strip enclosed by the edge f, the edge e extended beyond u, and the line parallel to e through w, is empty, and since $\mathrm{pc}(P)$ is connected we must therefore have that $\mathrm{pc}(P)$ is a subset of the closure of $\mathbb{H}^*_{\gamma(e),u} \cup \mathbb{H}^*_{\gamma(f),w}$. Now we reverse the order that edges e and f appear in the polygon, and let the new vertex introduced at their intersection be v'. Note that v' is convex, and that the operation has not made either u or w concave (this last statement follows from $|\gamma(e) - \gamma(f)| = 1$ and there being no two consecutive concave vertices in $\mathrm{pc}(P)$). Thus we have not increased the number of vertices and we have strictly decreased the number of concave vertices. While there are concave vertices left, repeat this edge reversing procedure.

The resulting polygon, Q, is a convex $\mathcal{S}$-polygon. We would like to show that $Q = \langle P \rangle$. Since $\langle P \rangle$ is by definition the intersection of all convex $\mathcal{S}$-polygons containing P, we just have to show that $Q \subset \langle P \rangle$. For this it is sufficient to have that whenever $P \subset \mathbb{H}^*_{u,a}$ it is also the case that $Q \subset \mathbb{H}^*_{u,a}$, for any $a \in \mathbb{R}^2$ and $u \in \mathcal{S}$. Now since $\mathrm{pc}(P) \subset \mathrm{conv}(P)$, we certainly have $P \subset \mathbb{H}^*_{u,a}$ implies $\mathrm{pc}(P) \subset \mathbb{H}^*_{u,a}$. Next, an easy induction shows us that each edge of Q contains a vertex from the original polygon P (but not necessarily at an endpoint), and we know each edge has outward normal vector belonging to $\mathcal{S}$; these observations together give us $Q \subset \langle P \rangle$ as required. Finally, to obtain $\langle P \rangle = Q$ from $\mathrm{pc}(P)$ we just changed the order of the sides around the polygon. $\square$

The power of Lemma 14 lies in the following inequality, which is the main subadditivity lemma.

Lemma 15. *Let D_1 and D_2 be droplets for $\mathcal{S}$ such that $D_1 \cap D_2 \neq \emptyset$ and let $u \in S^1$. Then*

$$\pi_u(\langle D_1 \cup D_2 \rangle) \leqslant \pi_u(D_1) + \pi_u(D_2).$$



$\langle D_1 \cup D_2 \rangle$
D_1
$\text{pc}(D_1 \cup D_2)$
D_2
$\pi_u(D_1)$
u
$\pi_u(D_2)$
$\pi_u(\langle D_1 \cup D_2 \rangle)$

Figure 5. Lemma 15 says that we always have the sum of the lengths of the two dashed projection line segments at most the length of the solid projection line segment.

Let P be a polygon in the plane, let $e_1, \ldots, e_m$ be its sides, and let $u \in S^1$. We define the *edgewise projection* $\pi'_u(P)$ of P in direction u by

$$\pi'_u(P) = \sum_{i=1}^{m} \pi_u(e_i).$$

Again, if D is a droplet for $\mathcal{S}$, we let $\pi'_u(D) := \pi'_u(\text{poly}(D))$. If P is convex then we have $\pi'_u(P) = 2\pi_u(P)$ for all directions u.

Proof of Lemma 15. We have the following inequalities.

$$\begin{aligned} 2\pi_u(D_1) + 2\pi_u(D_2) &= \pi'_u(D_1) + \pi'_u(D_2) \\ &\geqslant \pi'_u(D_1 \cup D_2) \\ &\geqslant \pi'_u(\text{pc}(D_1 \cup D_2)). \end{aligned}$$

The first inequality follows because the perimeter of $D_1 \cup D_2$ is contained within the perimeters of D_1 and D_2. The second inequality follows because $\text{pc}(D_1 \cup D_2)$ is obtained from $\operatorname{poly}(D_1 \cup D_2)$ by successively replacing piecewise linear segments of the boundary of $\operatorname{poly}(D_1 \cup D_2)$ by straight lines with the same end points, each such replacement preserving the inequality. By Lemma 14, $\operatorname{pc}(D_1 \cup D_2)$ and $\operatorname{poly}(\langle D_1 \cup D_2 \rangle)$ have the same edges, so we have

$$2\pi_u(D_1) + 2\pi_u(D_2) \geqslant \pi'_u(\langle D_1 \cup D_2 \rangle) = 2\pi_u(\langle D_1 \cup D_2 \rangle). \qquad \square$$

Lemma 14 also has the following corollary, which we do not use but we note for reference.

Corollary 16. *Let D_1 and D_2 be droplets for $\mathcal{S}$ such that $D_1 \cap D_2 \neq \emptyset$. Then*

$$p_E(\langle D_1 \cup D_2 \rangle) \leqslant p_E(D_1) + p_E(D_2). \qquad \square$$

Next we move on to describing the covering algorithm, which will be our approximation to the generalized bootstrap process. Let $\mathcal{U}$ be critical and let Λ be a droplet. Let $\hat{D}$ be a droplet that contains every update rule in $\mathcal{U}$. Choose a maximal set $\mathcal{B} = \{B_1, \dots, B_k\}$ of disjoint breakthrough blocks in A, and for each i, $1 \leqslant i \leqslant k$, let D_i^1 be a minimal droplet containing B_i. Suppose we have $D_1^t, \dots, D_{k_t}^t$. If there do not exist indices i and j and a site x such that

$$\operatorname{poly}(D_i^t) \cup \operatorname{poly}(D_j^t) \cup \operatorname{poly}(\hat{D} + x) \tag{8}$$

is connected, then stop. If there do exist such i, j and x, then choose one such triple, and let $D' = \langle D_i \cup D_j \rangle$. We construct $D_1^{t+1}, \dots, D_{k_{t+1}}^{t+1}$ as follows. From the list $D_1^t, \dots, D_{k_t}^t$, delete D_i^t and D_j^t and add D'. Repeat this procedure until there are no longer any droplets that can be combined. Let the final droplets be $D_1^\tau, \dots, D_{k_\tau}^\tau$. (We note that the order in which droplets are combined does not affect the final droplets, but the way in which the B_i are chosen may affect the final droplets.) If no matter how we choose $\mathcal{B}$ we always end with $k_\tau = 1$ and $E_1 = \Lambda$, we say that Λ is *strongly covered by A*, or when A is understood, that Λ is *strongly covered.*

The following series of lemmas collect together some information about the relationship between covered droplets and the generalized bootstrap process with initial set A.

Lemma 17. *Let $\mathcal{U}$ be critical and let Λ be a strongly covered droplet with diameter d. Then A contains $\Omega(d)$ disjoint breakthrough blocks.*

Proof. By Lemma 15, when droplets D_i^t and D_j^t combine under the strong covering algorithm we have the inequality

$$\pi_u(\langle D_i^t \cup D_j^t \rangle) \leqslant \pi_u(D_i^t) + \pi_u(D_j^t) + c_1,$$

where $c_1 = \pi_u(\hat{D})$ is a constant and u is an arbitrary unit vector. At each step of the algorithm we reduce the number of droplets by one, so there are $|\mathcal{B}| - 1$ steps in total. The droplet D_i^1 is contained in (a translate of) the droplet D that contains every update rule X in $\mathcal{U}$, so we have $\pi_u(D_i^1) \leqslant c_1$. Therefore, if Λ is strongly covered by A, we have

$$\operatorname{diam}(\Lambda) = \sup_{u \in S^1} \pi_u(\Lambda) \leqslant (2|\mathcal{B}| - 1)c_1,$$

so $|\mathcal{B}| = \Omega(d)$. $\square$

Lemma 18. *Let $\mathcal{U}$ be critical. Let Λ be a droplet, $\mathcal{B}$ a maximal set of breakthrough blocks in A, and $E_1, \dots, E_l$ the final droplets in the strong covering algorithm run on Λ with $\mathcal{B}$. Then $[A] \setminus A \subset E_1, \dots, E_l$.*

Proof. Let F be the set $A \setminus (E_1 \cup \cdots \cup E_l)$. The set $E_1 \cup \cdots \cup E_l \cup F$ is closed because each E_i is closed in itself, no two of the E_i are close enough to interact, and F does not contain any breakthrough blocks. The set $[A]$ is contained in $E_1 \cup \cdots \cup E_l \cup F$, since the latter is closed and contains A. Therefore

$$[A] \setminus A \subset (E_1 \cup \cdots \cup E_l \cup F) \backslash A \subset E_1 \cup \cdots \cup E_l. \qquad \square$$

Lemma 19. *Let $\mathcal{U}$ be critical and let Λ be a strongly covered droplet with diameter d. Then for any d' with $c_1 \leqslant d' \leqslant d/3$, Λ contains a strongly covered droplet D of diameter between d' and $3d'$.*

Proof. Lemma 15 gives

$$\pi_u(\langle D_1 \cup D_2 \rangle) \leqslant 3 \max\{\pi_u(D_1), \pi_u(D_2)\}$$

for all $u \in S^1$ whenever there exists x such that

$$\mathrm{poly}(D_1) \cup \mathrm{poly}(D_2) \cup \mathrm{poly}(\hat{D} + x)$$

is connected, provided $\max\{\pi_u(D_1), \pi_u(D_2)\}$ is at least $c_1 := \pi_u(\hat{D})$. Therefore

$$\begin{aligned} \mathrm{diam}(\langle D_1 \cup D_2 \rangle) &= \sup_{u \in S^1} \pi_u(\langle D_1 \cup D_2 \rangle) \\ &\leqslant \sup_{u \in S^1} 3 \max\{\pi_u(D_1), \pi_u(D_2)\} \\ &= 3 \max\{\mathrm{diam}(D_1), \mathrm{diam}(D_2)\}. \end{aligned}$$

So the maximum diameter over all the droplets at most triples at each step of the algorithm. $\square$

4.2. **Proof of Theorem 1.** In the introduction we dicussed the role of critical droplets in bootstrap percolation. 'Critical' in this context means 'about the right size such that the probability of there existing a droplet of this size that contains at least the minimal number of sites (or breakthrough blocks) needed to be internally spanned is minimized'. It turns out that 'about the right size' means 'about the logarithm of the diameter of the main droplet'. Before we turn to the proof of Theorem 1 we need two calculations. The first says approximately that the probability of there existing a strongly covered critical droplet is small. The second says approximately that the probability of there existing a strongly covered droplet containing the origin and having diameter no larger than $\log t$ is also small. The proof of Theorem 1 will then split into these two cases.

Before we begin, we define the main exponent, α_1. Let

$$\alpha_1 = \min\{|B| : B \text{ is a breakthrough block for some } X \in \mathcal{U}\}.$$

Lemma 20. *Let $\mathcal{U}$ be critical, let Λ be a droplet of diameter d, and let $c_1, c_2 > 0$ be constants. Let $p = p(d)$ satisfy*

$$p \leqslant \left(\frac{\lambda_1}{\log d}\right)^{\frac{1}{\alpha_1}}$$

for some sufficiently small constant $\lambda_1 > 0$. Finally, let G be the event that there exists a droplet $D \subset \Lambda$ with diameter between $c_2 \log d$ and $3c_2 \log d$ such that $D \cap \Pi_p$ contains at least $c_1 \log d$ breakthrough blocks. Then there exists a constant $c > 0$ such that

$$\mathbb{P}_p(G) \leqslant \exp(-c \log d).$$

Proof. The number of droplets contained in Λ with diameter between $c_2 \log d$ and $3c_2 \log d$ is at most

$$|\Lambda|(c_3 c_2 \log d)^s \leqslant c_4 d^2 (\log d)^s,$$

for some positive constants c_4 and c_5. A droplet of diameter at most $3c_2 \log d$ contains at most $c_5 (\log d)^2$ breakthrough blocks, for some c_5. Recall that each breakthrough block has size at least α_1. The probability that there exists a droplet D of diameter at most $3\log d$ such that $D \cap \Pi_p$ contains at least $c_1 \log d$ disjoint breakthrough blocks is therefore at most

$$c_4 d^2 (\log d)^s \binom{c_5 (\log d)^2}{c_1 \log d} p^{\alpha_1 c_1 \log d}.$$

By Stirling's formula, this is at most

$$c_4 d^2 (\log d)^s \left(\frac{c_5 (\log d)^2 e}{c_1 \log d} \right)^{c_1 \log d} p^{\alpha_1 c_1 \log d},$$

so we have

$$\mathbb{P}_p(G) \leqslant \exp\Big(\log c_4 + 2\log d + s \log\log d + \\ c_1 \log d \big(\log(c_5 e / c_1) + \log\log d \big) - c_1 (\log d) \alpha_1 \log(1/p) \Big).$$

Now, since $\alpha_1 \log(1/p) \geqslant \log\log d - \log \lambda_1$, it follows that

$$\mathbb{P}_p(G) \leqslant \exp\big(c_6 + s \log\log d + (c_7 + c_1 \log \lambda_1) \log d \big) \tag{9}$$

for some constants c_6 and c_7. For small enough λ_1 the coefficient of $\log d$ inside the exponential is negative, so for large enough d and some $c > 0$, we have

$$\mathbb{P}_p(G) \leqslant \exp(-c \log d). \qquad \square$$

Lemma 21. *Let $\mathcal{U}$ be critical and let $A = \Pi_p$. Let H be the event that the origin is contained in a strongly covered droplet with diameter at most $\log t$. Then there exists a constant $c' > 0$ such that*

$$\mathbb{P}_p(H) \leqslant \exp(-c' \log d).$$

Proof. Let H_i be the event that the strongly covered droplet D containing the origin has diameter between $i-1$ and i. There are at most $c_1 i^s i^2$ droplets of diameter between $i-1$ and i that contain 0, and each of them contains at most $c_2 i^2$ breakthrough blocks. Furthermore, by Lemma 17, $D \cap \Pi_p$ contains at least $c_3 i$ disjoint breakthrough blocks. Let $d = \operatorname{diam}(D(\log t))$, so $d \leqslant c_4 \log t$ for some c_4. Putting these together, we have

$$\mathbb{P}_p(H) \leqslant \sum_{i=1}^{d} \mathbb{P}_p(H_i) \leqslant \sum_{i=1}^{d} c_1 i^{s+2} \binom{c_2 i^2}{c_3 i} p^{\alpha_1 c_3 i} \leqslant c_1 \sum_{i=1}^{d} i^{s+2} \left(\frac{c_2 i e p^{\alpha_1}}{c_3} \right)^{c_3 i}.$$

There exists a constant c_5 such that $c_1 i^{s+1} \leqslant c_5 e^{c_3 i}$, so

$$\mathbb{P}_p(H) \leqslant c_5 \sum_{i=1}^{d} \left(\frac{c_2 i e^2 p^{\alpha_1}}{c_3} \right)^{c_3 i} \leqslant c_5 \sum_{i=1}^{d} \left(\frac{c_2 i e^2 \lambda_1}{c_3 \log t} \right)^{c_3 i} \leqslant c_5 \sum_{i=1}^{d} \left(\frac{c_2 i e^2 \lambda_1 c_4}{c_3 d} \right)^{c_3 i}.$$

Letting $\lambda_1' = c_2 e^2 \lambda_1 c_4 / c_3$ be an arbitrarily small constant, we can write this as

$$\mathbb{P}_p(H) \leqslant c_5 \sum_{i=1}^{d} \left(\frac{\lambda_1' i}{d} \right)^{c_3 i}.$$

We then have

$$\begin{aligned}\mathbb{P}_p(H)c_5^{-1} &\leqslant \sum_{i=1}^{\sqrt{d}} \left(\frac{\lambda_1' i}{d}\right)^{c_3 i} + \sum_{i=\sqrt{d}}^{d} \left(\frac{\lambda_1' i}{d}\right)^{c_3 i} \\ &\leqslant \sum_{i=1}^{\infty} \left(\frac{\lambda_1' \sqrt{d}}{d}\right)^{c_3 i} + \sum_{i=\sqrt{d}}^{\infty} (\lambda_1')^{c_3 i} \\ &= \left(\frac{\lambda_1'}{\sqrt{d}}\right)^{c_3} \frac{1}{1-(\lambda_1'/\sqrt{d})^{c_3}} + (\lambda_1')^{c_3\sqrt{d}} \frac{1}{1-\lambda_1'} \\ &\leqslant c_6 \exp(-c_7 \log d)\end{aligned}$$

for some constants c_6 and c_7, as required. □

Finally we are ready to prove Theorem 1. The argument runs as follows. First we observe that the event that the origin is infected by time t does not depend on anything that happens outside of a droplet centred on the origin of size proportional to t. Next, we show that either the origin is strongly covered inside a droplet of diameter $\log t$, or there exists a strongly covered critical droplet not too far from the origin. We know that the probability of the first of these is small from Lemma 21, and that the probability of the second of these is also small from Lemma 20.

Proof of Theorem 1. Let E_t be the event the that origin is infected by time t. Given k, let $D(k)$ be the minimal droplet with the property that the Euclidean distance between every point on $\operatorname{peri}(D(k))$ and the origin is at least k; this is well-defined since we can consider sides independently. There exists a constant c_1 such that if $T = c_1 t$, then E_t implies the event $F = \{0 \in [D(T) \cap A]\}$. For example, we could take

$$c_1 = \max\{\|x\| : x \in X, X \in \mathcal{U}\}.$$

(We think of c_1 as being the maximum speed information can travel.)

Let G be the event that there exists a droplet in $D(T)$ of diameter between $\log t$ and $3 \log t$ that contains at least $c_2 \log t$ disjoint breakthrough blocks, where c_2 is the implicit constant in the second part of Lemma 17. Let H be the event that the origin is contained in a strongly covered droplet which is itself contained in $D(\log t)$.

If F occurs and $0 \notin A$, then by Lemma 18 the origin lies inside a strongly covered droplet $D \subset D(T)$. If H^c also occurs, D cannot be contained entirely within $D(\log t)$, so it must have diameter between $\log t$ and $2T$. Lemma 19 tells us that D contains a strongly covered droplet D' with diameter between $\log t$ and $3 \log t$, so $F \cap \{0 \notin A\} \cap H^c$ implies G, or equivalently, F implies $G \cup H \cup \{0 \in A\}$. Thus,

$$\mathbb{P}_p(E) \leqslant \mathbb{P}_p(F) \leqslant \mathbb{P}_p(G) + \mathbb{P}_p(H) + \mathbb{P}_p(0 \in A). \tag{10}$$

Finally we show that each of the probabilities on the right of (10) is $o(1)$. First, $\mathbb{P}_p(0 \in A) = p = o(1)$. Next, by applying Lemma 20 to $\Lambda = D(T)$ (which has diameter $O(t)$), we have $c_3 > 0$ such that for large enough t,

$$\mathbb{P}_p(G) \leqslant \exp(-c_3 \log t) \to 0,$$

and similarly, Lemma 21 gives immediately some $c_4 > 0$ such that

$$\mathbb{P}_p(H) \leqslant \exp(-c_4 \log t) \to 0,$$

which completes the proof. □

4.3. **Quasi-stability.** We now start building up to the proof of the upper bound, Theorem 2.

Given a stable direction u, we define ℓ_u^l to be the set of those points x in ℓ_u such that the origin lies to the left of x as one looks in the direction of u. Analogously, we define ℓ_u^r to be the set of those points x in ℓ_u such that the origin lies to the right of x as one looks in the direction of u. A *u-left-block* is a breakthrough block $B = X \cap \mathbb{H}_u^c$ (for some $X \in \mathcal{U}$) such that $B \subset \ell_u^l$. A *u-right-block* is a breakthrough block $B = X \cap \mathbb{H}_u^c$ (for some $X \in \mathcal{U}$) such that $B \subset \ell_u^r$. These special breakthrough blocks are so-called because we think of them as being the means of growing left or right along a side of a droplet.

Lemma 22. *Let $\mathcal{U}$ be critical. Then there exists a u-left-block and a u-right-block for every stable direction u.*

Proof. Let $u = u(\theta)$ be stable. Since $\mathcal{U}$ is critical, there exists an update rule X and $\delta > 0$ such that $u(\phi)$ is destabilized by X for every ϕ in the open interval $I = (\theta, \theta + \delta)$. Thus

$$X \subset \bigcap_{\phi \in I} \mathbb{H}_{u(\phi)} \subset \mathbb{H}_u \cup \ell_u^l,$$

so $X \cap (\mathbb{H}_u)^c$ is a u-left-block. u-right-blocks exist by symmetry. ☐

Given a breakthrough block B, let $\|B\|$ be the *length* of B, defined by

$$\|B\| := \max\{\|x\| \,:\, x \in B\}.$$

For each stable direction u, let $\mathcal{B}_u^l$ be the set of all u-left-blocks. Let $B_u^l = X_u^l \cap \mathbb{H}_u$ be a u-left-block with minimal length, so if $B \in \mathcal{B}_u^l$ then $\|B_u^l\| \leqslant \|B\|$. We call B_u^r *the u-left-block* and X_u^l *the u-left-rule*. Similarly, let $\mathcal{B}_u^r$ be the set of all u-right-blocks and let $B_u^r = X_u^r \cap \mathbb{H}_u$ be a u-right-block with minimal length, so if $B \in \mathcal{B}_u^r$ then $\|B_u^r\| \leqslant \|B\|$. We call B_u^r *the u-right-block* and X_u^r *the u-right-rule*.

These definitions allow us to define α_2, one of the two constants in Theorem 2.

$$\alpha_2 := \max\big\{\{\|B_u^l\| \,:\, u \in \mathcal{S}\} \cup \{\|B_u^r\| \,:\, u \in \mathcal{S}\}\big\}. \tag{11}$$

Quasi-stable directions. The need for quasi-stable directions exists because without them we do not have a basis for growing droplets into droplets. Before we give the definition, we illustrate the situation with an example. Suppose X_1 and X_2 are two update rules such that $\Theta(X_i) = (\phi_i, \psi_i)$ for $i = 1, 2$, and

$$\phi_1 < \phi_2 < \psi_1 < \psi_2.$$

Thus, there are no stable directions between $u(\phi_1)$ and $u(\psi_2)$. Suppose (although it does not much matter), that in fact $u(\phi_1)$ and $u(\psi_2)$ are stable directions, so droplets for $\mathcal{S}$ have consecutive sides with outward normals in the directions of $u(\phi_1)$ and $u(\psi_2)$. Now suppose we are growing leftwards along the $u(\phi_1)$ side, and we are nearly at the corner of the droplet. How are we going to reach the corner? X_1 does not fit inside the droplet near the corner because $\psi_1 < \psi_2$, so it partly lies outside the $u(\psi_2)$ side. X_2 does not fit inside the droplet near the corner either, because $\phi_2 > \phi_1$, so it partly lies outside the $u(\phi_1)$ side (see Figure 6). So there is no rule that enables us to reach the corner, and we are stuck. The solution is to introduce a new direction u, which we shall treat as though it were a stable direction, and which is such that $\phi_2 < \theta(u) < \psi_1$ (and otherwise arbitrary). If we add u to $\mathcal{S}$, we are now able to grow to the corners of droplets for $\mathcal{S}$ (or at least, we are able to grow to the corners adjacent to the u sides of droplets).

Let $\mathcal{S} = \{u_1, \dots, u_s\}$, where the indexing is such that for each r mod s, $u_r = u(\theta_r)$ and $u_{r+1} = u(\theta_{r+1})$ are consecutive stable directions going anticlockwise around S^1. Fix r and let $X_{i_1}, \dots, X_{i_{k_r}}$ be a minimal collection of update rules such that



FIGURE 6. An illustration of the need for quasi-stable directions. The shaded area enclosed by the dashed lines is $\mathbb{H}_{u(\phi_1)} \cap \mathbb{H}_{u(\psi_1)}$, which contains the rule X_1. The shaded area enclosed by the dotted lines is $\mathbb{H}_{u(\phi_2)} \cap \mathbb{H}_{u(\psi_2)}$, which contains the rule X_2. A corner of a typical droplet for $\mathcal{S}$ would look like the dark grey shaded area, which contains neither X_1 nor X_2, so neither rule can reach the corner. There is a slight abuse of notation in the labelling: the directions should be $u(\phi_1)$, etc., not ϕ_1, etc.

(i) $X_{i_1} = X^l_{u_r}$;
(ii) $X_{i_{k_r}} = X^r_{u_{r+1}}$;
(iii) $\theta^-(X_{i_{j+1}}) < \theta^+(X_{i_j})$ for all $1 \leqslant j < k_r$.

Such a collection exists because $u(\theta)$ is not stable for any $\theta \in (\theta_r, \theta_{r+1})$. For each $1 \leqslant j < k_r$ choose an arbitrary

$$\phi_j \in (\theta^-(X_{i_{j+1}}), \theta^+(X_{i_j})) \tag{12}$$

such that $u(\phi_j)$ has rational slope, and let $Q_r := \{u(\phi_j) \,:\, 1 \leqslant j < k_r\}$. Q_r will be empty if $X^l_{u_r} = X^r_{u_{r+1}}$, since in that case the minimal collection of rules is just $\{X^l_{u_r}\}$. We define the *quasi-stable set* $\mathcal{S}_1$ to be

$$\mathcal{S}_1 := \mathcal{S} \cup \bigcup_{r=1}^{s} Q_r.$$

Each $\theta \in \mathcal{S}_1 \setminus \mathcal{S}$ is called a *quasi-stable direction.* We re-index the u_i so that $\mathcal{S}_1 = \{u_1, \ldots, u_{s_1}\}$, with the directions indexed in order anticlockwise around S^1 (starting at $u(0)$, say), and the index always taken mod s_1. A *quasi-droplet* is droplet for $\mathcal{S}_1$.

If $u \in \mathcal{S}_1 \setminus \mathcal{S}$ we define $X^l_u = X^r_u = B^l_u = B^r_u = \emptyset$. Intuitively this is because u is not a stable direction, so the empty update $X = \emptyset$ is enough to grow beyond $\mathbb{H}_u$.

The next lemma ensures that update rules fit into the the corners of quasi-droplets, which will also mean that we can grow to the corners of quasi-droplets. It is the key property of the quasi-stable set.

Lemma 23. *Let $\mathcal{U}$ be critical. Then there exists a positive integer H such that*

$$X^l_{u_i} \subset (B^l_{u_i} \cup \mathbb{H}_{u_i}) \cap \mathbb{H}_{u_{i+1}}(1) \cap \mathbb{H}_{u_{i-1}}(H) \cap \mathbb{H}_{-u_i}(H)$$

and

$$X^r_{u_i} \subset (B^r_{u_i} \cup \mathbb{H}_{u_i}) \cap \mathbb{H}_{u_{i-1}}(1) \cap \mathbb{H}_{u_{i+1}}(H) \cap \mathbb{H}_{-u_i}(H)$$

for each $1 \leqslant i \leqslant s_1$.

Proof. Fix i. If $u_i \in \mathcal{S}_1 \setminus \mathcal{S}$ then the assertion is trivial because the u-left-rule and the u-right-rule are both the empty set. So suppose $u_i \in \mathcal{S}$. Clearly

$$X^l_{u_i} \subset \mathbb{H}_{-u_i}(H_i) \cap \mathbb{H}_{u_{i-1}}(H_i) \qquad \text{and} \qquad X^r_{u_i} \subset \mathbb{H}_{-u_i}(H_i) \cap \mathbb{H}_{u_{i+1}}(H_i)$$

for some H_i, since update rules are bounded, so we set $H = \max_i H_i$. Also, $X^l_{u_i} \subset B^l_{u_i} \cup \mathbb{H}_{u_i}$ because $X^l_{u_i}$ is the u-left-rule and therefore $X^l_{u_i} \setminus B^l_{u_i} \subset \mathbb{H}_{u_i}$. It remains to show that $X^l_{u_i} \subset \mathbb{H}_{u_{i+1}}(1)$. If this doesn't hold then $\theta^+(X^l_{u_i}) < \theta(u_{i+1})$. If u_{i+1} belongs to $\mathcal{S}_1 \backslash \mathcal{S}$ then this is a contradiction by (12). If u_{i+1} belongs to $\mathcal{S}$ then this is also a contradiction because we would have $\theta^+(X^l_{u_i}) = \theta(u_{i+1})$. □



FIGURE 7. Lemma 23 says that $X^l_{u_i}$ is contained in something like the shaded area here, excluding sites on the dashed lines, but including the breakthrough block $B^l_{u_i}$, shown as the sites inside the solid reectangle.

4.4. **Growth of quasi-droplets.** In the previous section we formally introduced quasi-stability, and proved in Lemma 23 that with the addition of quasi-stable directions to the stable set, we are able to reach the corners of quasi-droplets. Our task in this section is to describe the growth of quasi-droplets.

Let D be a quasi-droplet. For each quasi-stable direction u set $i_u(D)$ to be the unique i such that

$$\ell_u(i) \cap D \neq \emptyset \qquad \text{and} \qquad \ell_u(i+1) \cap D = \emptyset.$$

Thus, D can be written as

$$D = \bigcap_{u \in \mathcal{S}_1} \mathbb{H}_u(i_u(D) + 1).$$

In the next lemma we prove the existence of a nested sequence of quasi-droplets with linearly increasing side lengths for any given initial quasi-droplet D. The subtlety of the lemma is merely that we are working on a discrete grid rather than $\mathbb{R}^2$. We expect the vertices of the nested sequence to lie on or near the angle bisectors of the first quasi-droplet, and most of the proof is concerned with showing that this is indeed possible. Once we have the sequence of quasi-droplets, the remaining results of this section ensure that we can think of growth of quasi-droplets in terms of this sequence, and not have to worry about shrinking sides.

Lemma 24. *Let $\mathcal{U}$ be critical. Then there exist positive integers $d_1, \dots, d_{s_1}$ such that if D is any quasi-droplet then the sequence of quasi-droplets $(D_k)_{k=1}^{\infty}$ given by*

$$D_k = \bigcap_{i=1}^{s_1} \mathbb{H}_{u_i}(i_{u_i}(D) + (k-1)d_i + 1) \tag{13}$$

have the following properties for every $k \geqslant 1$:

(i) $D_k \subset D_{k+1}$*;*
(ii) $\|\operatorname{side}_i(D_k)\|$ *increases linearly in k for each fixed $i \leqslant s_1$;*

Proof. Our main task will be finding a suitable droplet D_2 (by definition, $D_1 = D$) with corresponding positive integers $d_1, \ldots, d_{s_1}$ that do not depend on D. Verifying properties (i)-(ii) for the resulting sequence of droplets $(D_k)_{k=1}^{\infty}$ will follow with little extra work.

Let u_i be a quasi-stable direction and let v_i and v_{i+1} be the end vertices of the side of peri(D) corresponding to u_i with v_i to the right of v_{i+1} as one looks in the direction of u_i. Let b_i and b_{i+1} be angle bisectors of peri(D) at v_i and v_{i+1} respectively. Let $\mu_i(j)$ be the subset of $\ell_{u_i}(i_{u_i}(D) + j)$ consisting of sites on or between the bisectors b_i and b_{i+1} for each $j \geqslant 0$ (so $\mu_i(0)$ is the u_i side of D), and analogously let $\mu_i^*(j)$ be the closed line segment of $\ell_{u_i}^*(i_{u_i}(D) + j)$ consisting of all points on or between the bisectors b_i and b_{i+1}. The Euclidean length of $\mu_i^*(j)$ increases linearly in j, and moreover the rate of increase depends only on i and $\mathcal{S}_1$, not on D. Accordingly, let $m_i > 0$ be such that

$$\|\mu_i^*(j)\| = l_i + m_i j, \tag{14}$$

where l_i is the Euclidean length of the u_i side of peri(D) and m_i does not depend on D.

Let y_i^- be the distance between the point of intersection of $\ell_{u_{i-1}}^*(i_{u_{i-1}}(D))$ and $\mu_i^*(1)$, and the point of intersection of $\mu_i^*(1)$ and b_i. Let y_i^+ be the distance between the point of intersection of $\ell_{u_{i+1}}^*(i_{u_{i+1}}(D))$ and $\mu_i^*(1)$, and the point of intersection of $\mu_i^*(1)$ and b_{i+1}.

Let d_i' be the minimal positive integer such that

$$m_i d_i' \geqslant y_i^- + y_i^+ + 1. \tag{15}$$

Note that d_i' depends only on m_i, y_i^- and y_i^+, none of which depend on D. Let h_i be the perpendicular distance between $\mu_i^*(0)$ and $\mu_i^*(d_i')$, and let $h = \max\{h_i : 1 \leqslant i \leqslant s_1\}$. Finally we are in a position to define d_i. We define d_i to be the minimal positive integer such that the perpendicular distance between $\mu_i^*(0)$ and $\mu_i^*(d_i)$ is at least h.

We need to check properties (i) and (ii) for the D_k as defined in (13) for each $k \geqslant 1$. (i) is trivial by construction for every $k \geqslant 1$, so we just have to check the linearity property. For each $1 \leqslant i \leqslant s_1$ and $k \geqslant 0$ let $\nu_i(k) := \ell_{u_i}^*(i_{u_i}(D) + kd_i)$ and let $w_i(k)$ be the point of intersection of $\nu_{i-1}(k)$ with $\nu_i(k)$. Let the intersections of $\nu_{i-1}(1)$ and $\nu_i(1)$ with b_i occur at ξ_i^- and ξ_i^+ respectively (see Figure 8).

Suppose $w_i(1)$ lies to the left (i.e. on the u_i side) of b_i. This means ξ_i^- is closer to v_i than ξ_i^+ is to v_i. We claim the distance between $w_i(1)$ and ξ_i^+ is at most y_i^-. If this were not the case, $\mu_i^*(d_i - 1)$ would also intersect $\nu_{i-1}(1)$ on the u_i side of b_i, so since b_i is a bisector we would have the perpendicular distance of $\mu_i^*(d_i - 1)$ from $\mu_i^*(0)$ also at least h, contradicting the definition of d_i.

Similarly, suppose $w_{i+1}(1)$ lies to the right (i.e. on the u_i side) of b_{i+1}. This means ξ_{i+1}^+ is closer to v_{i+1} than ξ_{i+1}^- is to v_{i+1}. We claim the distance between $w_{i+1}(1)$ and ξ_{i+1}^- is at most y_i^+. If this were not the case, $\mu_i^*(d_i - 1)$ would also intersect $\nu_{i+1}(1)$ on the u_i side of b_{i+1}, so since b_{i+1} is a bisector we would have the perpendicular distance of $\mu_i^*(d_i - 1)$ from $\mu_i^*(0)$ also at least h, contradicting the definition of d_i.

It follows from these last two paragraphs, together with (14) and (15) that

$$\|\operatorname{side}_i(D_2)\| \geqslant \|\mu_i^*(d_i)\| - (y_i^- + y_i^+) \geqslant l_i + m_i d_i - (y_i^- + y_i^+) \geqslant l_1 + 1.$$

The points $w_i(0), w_i(1), w_i(2), \ldots$ lie equally spaced on a straight line, so the difference $\|\operatorname{side}_i(D_{k+1})\| - \|\operatorname{side}_i(D_k)\|$ does not vary with k. When $k = 1$ the difference is positive (in fact it is at least 1), so the lengths increase linearly in k, which is property (ii). $\square$

Lemma 25. *Let $\mathcal{U}$ be critical. Then there exists a constant J such that the following holds. Let D be a quasi-droplet all of whose sides have lattice length at least $k + 2J$. Then*

$$\left|\ell_{u_i}(i_{u_i}(D) + d_i) \cap \mathbb{H}_{u_{i-1}}(i_{u_{i-1}}(D) + 1) \cap \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D) + 1)\right| \geqslant k. \tag{16}$$



FIGURE 8. Lemma 24 with $d_{i-1} = 3$, $d_i = 2$ and $d_{i+1} = 5$.

Proof. This is straightforward. Essentially the lemma is equivalent to the statement that the rate at which sides of D shrink is a constant that depends only on $\mathcal{S}_1$.

For each $i \leqslant s_1$, let l_i be the (Euclidean) length of the u_i side of D, let l_i' be the length of the line segment of $\ell^*_{u_i}(i_{u_i}(D) + d_i)$ lying between $\ell^*_{u_{i-1}}(i_{u_{i-1}}(D))$ and $\ell^*_{u_{i+1}}(i_{u_{i+1}}(D))$, and let ϕ_i be such that $\cos\phi_i = \langle u_i, u_{i+1}\rangle$. Recalling that η_u is the gap between lines of direction u, we have

$$l_i - l_i' = d_i\eta_{u_i}(\cot\phi_i + \cot\phi_{i-1}).$$

Now recalling that δ_u is the gap between consecutive sites on a line of direction u, a segment of a line of direction u of this length contains at most

$$J_i := \lfloor d_i\eta_{u_i}(\cot\phi_i + \cot\phi_{i-1})/\delta_{u_i}\rfloor + 1$$

sites, which does not depend on D. Therefore we can take $J = \max_i J_i$. □

Let K be a constant and let D be a quasi-droplet such every side of D contains at least K sites. Fix i and let ξ be the line segment given by

$$\xi := \ell_{u_i}(i_{u_i}(D) + 1) \cap \mathbb{H}_{u_{i-1}}(i_{u_{i-1}}(D) + 1) \cap \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D) + 1).$$

Let ρ be all the sites in ξ excluding the left-most H sites and the right-most H sites. We choose K large enough so that $|\rho|$ is at least α_2.

We are now in a position to prove our first result about the growth of quasi-droplets. We show that the presence of a breakthrough block on the side of a (sufficiently large) quasi-droplet results in infection spreading along the side to the corners.

Lemma 26. *Let $\mathcal{U}$ be critical. Let D, ξ and ρ be as above, let σ be any set of α_2 consecutive sites in ρ, and let K be sufficiently large. Then*

$$D \cup \xi \subset [D \cup \sigma]. \tag{17}$$

Proof. We prove (17) one site at a time, starting with the first site in ξ immediately to the left of σ (as one looks in the direction of u_i), then continuing inductively to the boundary, and then doing the same on the right of σ. Let $x(l)$ be the lth site in ξ to the left of σ, and let $y(l)$ be the lth site in ξ to the right of σ.

Starting the induction, let

$$D(l) := D \cup \sigma \cup \{x(1), \dots, x(l-1)\}.$$

We must show $X^l_{u_i} + x(l) \subset D(l)$. By Lemma 23,

$$X^l_{u_i} \subset (B^l_{u_i} \cup \mathbb{H}_{u_i}) \cap \mathbb{H}_{u_{i+1}}(1) \cap \mathbb{H}_{u_{i-1}}(H) \cap \mathbb{H}_{-u_i}(H),$$

so it is enough to show that the right-hand side of this equation, translated by $x(l)$, is a subset of $D(l)$. First, we have $B^l_{u_i} + x(l) \subset D(l)$, because the latter set contains at least α_2 sites to the right of $x(l)$ in ξ, and by the definitions of a u_i-left-rule and α_2, $B^l_{u_i}$ is a subset of $\ell^l_{u_i}$ such that $\|B^l_{u_i}\| \leqslant \alpha_2$. Therefore we just have to show

$$\big(\mathbb{H}_{u_i} \cap \mathbb{H}_{u_{i+1}}(1) \cap \mathbb{H}_{u_{i-1}}(H) \cap \mathbb{H}_{-u_i}(H)\big) + x(l) \subset D(l).$$

The left-hand side of this equation is contained in the quasi-droplet

$$\tilde{D}(l) := \left(\mathbb{H}_{u_i} \cap \mathbb{H}_{u_{i+1}}(1) \cap \bigcap_{i'=i+1}^{i-2} \mathbb{H}_{u_{i'}}(c_1) \cap \mathbb{H}_{u_{i-1}}(H)\right) + x(l)$$

for some constant c_1 (which does not depend on i or whether we are moving left or right), and the right-hand side contains the quasi-droplet D, so it suffices to show $\tilde{D}(l) \subset D$, or equivalently,

$$i_{u_{i'}}(\tilde{D}(l)) \leqslant i_{u_{i'}}(D) \tag{18}$$

for all $1 \leqslant i' \leqslant s_1$. When $i' = i$ we have this automatically. When $i' = i+1$ we have (18) because $\xi \subset \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D)+1)$ by the definition of ξ, and the boundary of the half-plane here is the u_{i+1} boundary of D. When $i' = i-1$, we know by the definition of ρ that there are at least H sites to the right of $x(l)$ in ξ, all contained in distinct lines of direction u_{i-1}, so we have (18). For $i' \notin \{i-1, i, i+1\}$, (18) holds simply by taking K, the minimum side length of D, sufficiently large.

We have proved that the $x(l)$ all belong to $[D \cup \sigma]$. The result for $y(l)$ now follows by symmetry. □

Let D be a quasi-droplet all of whose sides contain at least K sites, where K is to be determined, but is certainly at least large enough for Lemma 26 to hold. Let $(D_k)_{k=1}^{\infty}$ be the sequence of droplets given by Lemma 24. Let $\rho(i,0,k)$ be the elements of the u_i side of D_k excluding the left-most H points and the right-0most H points. We define $\rho(i,j,k)$ for $1 \leqslant j \leqslant d_i$ to be the set of sites in

$$\ell_{u_i}(i_{u_i}(D_k)+j) \cap \mathbb{H}_{u_{i-1}}(i_{u_{i-1}}(D_k)+1) \cap \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D_k)+1)$$

excluding the left-most and right-most H sites.

Lemma 27. *Let $\mathcal{U}$ be critical. Let D be as above and let K be sufficiently large (as a function of $\mathcal{S}_1$ only). For each triple i,j,k let $\sigma(i,j,k)$ be any set of α_2 consecutive sites in $\rho(i,j,k)$. Then for each $k \geqslant 1$ we have*

$$D_{k+1} \subset \Big[D_k \cup \bigcup_{i=1}^{s_1} \bigcup_{j=1}^{d_i} \sigma(i,j,k)\Big]. \tag{19}$$

Proof. Let $\Delta_k = D_{k+1} \setminus D_k$. We divide Δ_k up into the following line segments:

$$\begin{aligned}
\xi(1,j,k) &:= \ell_{u_1}(i_{u_1}(D_k)+j) \cap \mathbb{H}_{u_{s_1}}(i_{u_{s_1}}(D_k)+1) \cap \mathbb{H}_{u_2}(i_{u_2}(D_k)+1);\\
\xi(i,j,k) &:= \ell_{u_i}(i_{u_i}(D_k)+j) \cap \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D_k)+1) \cap \Delta_k;\\
\xi(s_1,j,k) &:= \ell_{u_{s_1}}(i_{u_{s_1}}(D_k)+j) \cap \Delta_k.
\end{aligned}$$

Observe that with these definitions we have

$$\bigcup_{i=1}^{s_1} \bigcup_{j=1}^{d_i} \xi(i,j,k) = \Delta_k. \tag{20}$$

For each $k \geqslant 1$, we prove (19) one line segment at a time, starting with $i = j = 1$ and increasing j until we reach $j = d_1$ with $i = 1$ fixed, then moving on to $i = 2$ and $j = 1$

and again continuing until we reach $j = d_2$ with $i = 2$ fixed, and so on. We prove (19) for each segment $\xi(i, j, k)$ using Lemma 26.

Starting then with $i = 1$, let

$$D(1, j, k) := D_k \cup \bigcup_{j'=1}^{j-1} \xi(i, j', k),$$

and notice that $D(1, j, k)$ is a quasi-droplet. In order to use Lemma 26, we need a lower bound on the number of sites on the u_1 side of $D(1, j, k)$. By Lemma 25, this number is at least the number of sites on the u_1 side of D minus a constant, for all j and k. Provided K is large enough, it follows from Lemma 26 applied to $D(1, j, k)$ and $\sigma = \sigma(1, j, k)$ that, inductively, $\xi(1, 1, k), \ldots, \xi(1, d_1, k)$ are all contained in

$$\Big[D_k \cup \bigcup_{i=1}^{s_1} \bigcup_{j=1}^{d_i} \sigma(i, j, k)\Big]. \tag{21}$$

For general i, let

$$D(i, j, k) := D_k \cup \bigcup_{i'=1}^{i-1} \bigcup_{j'=1}^{d_i} \xi(i', j', k) \cup \bigcup_{j'=1}^{j-1} \xi(i, j', k).$$

Again we observe that $D(i, j, k)$ is a quasi-droplet for every triple i, j, k. Furthermore, the number of sites along each side is greater here than it was in the $i = 1$ case, so the same argument as we used there easily gives us that Lemma 26 is applicable. Therefore, inductively, we have $\xi(2, 1, k), \ldots, \xi(2, d_2, k), \ldots, \xi(s_1, 1, k), \ldots, \xi(s_1, d_{s_1}, k)$ all contained in the set (21), as required. □

4.5. **Proof of Theorem 2.** The proof of Theorem 2 has the same structure as proofs of other theorems of this type (for example, Aizenman and Lebowitz [1] and Holroyd [16]). We divide the area surrounding the origin into disjoint squares, each of side length roughly $(\log t)^3$. We then prove that, with high probability, at least one of the squares contains an internally spanned critical droplet. Finally we show that every sufficiently long line segment of sites, in directions parallel to the sides of quasi-droplets, contains a breakthrough block. These breakthrough blocks enable the critical droplet to grow at least as far as the origin, completing the proof.

In the lower bound we were able to show that if a strongly covered droplet had a large projection in any single direction, then it had to contain a large number of breakthrough blocks. Thus, the definition of 'diameter' was one-dimensional: we only required the droplet to have one large projection; so for example, long thin rectangles would still have large diameter. Here in the upper bound setting, the one-dimensional definition is not enough: we rely heavily on the capacity to grow in all directions at once. Thus we need new (albeit related) notions of the size of the droplet, such that their being large guarantees that the droplet is large in a two-dimensional sense.

Given a quasi-droplet D, let $\phi(D)$ and $\Phi(D)$ denote, respectively, the minimum and maximum number of sites on sides of D (these being taken over all quasi-stable directions), and let $\phi^*(D)$ and $\Phi^*(D)$ denote, respectively, the minimum and maximum Euclidean side lengths of D. There exist positive constants such that $\phi(D) \leqslant c\phi^*(D) \leqslant C\phi(D)$ and $\Phi(D) \leqslant c\Phi^*(D) \leqslant C\Phi(D)$. Given any d, it is clear that there exists a quasi-droplet D with $\Phi^*(D) \leqslant d \leqslant s_1\phi^*(D)$, and therefore $\Phi(D) \leqslant cd \leqslant Cs_1\phi(D)$. We call quasi-droplets for which there exists d such that these inequalities hold *canonical quasi-droplets.* (The parameter d is not necessarily the diameter of these canonical quasi-droplets, but it is within a constant factor of the diameter.)

Before we prove Theorem 2, we need one tool from probability theory. An event $A \subset \{0,1\}^{\mathbb{Z}^2}$ is said to be *increasing* if whenever $\omega \in A$ and $\omega' \geqslant \omega$ under the product order (so all infected sites in ω are infected in ω'), it follows that $\omega' \in A$. The lemma that we need was first proved by Harris [15] in 1960, who used it in his work on bond percolation in the plane. Harris's Lemma is a special case of an inequality due to Fortuin, Kasteleyn and Ginibre [9], widely known as the FKG inequality.

Lemma 28 (Harris's Lemma [15])**.** *Let A and B be increasing events. Then*

$$\mathbb{P}_p(A \cap B) \geqslant \mathbb{P}_p(A)\mathbb{P}_p(B).$$

The first step in the proof of Theorem 2 is a lower bound on the probability that a small internally spanned droplet continues growing indefinitely. The proof is a generalization of van Enter's proof [19] that the critical probability for bootstrap percolation on the plane is zero.

Lemma 29. *Let $\mathcal{U}$ be critical. Let D be a quasi-droplet all of whose sides have lattice length at least K, and let $(D_k)_{k=1}^{\infty}$ be the sequence of droplets given by Lemma 24. Then there exists a constant $\lambda_1 > 0$ such that we have the following bound, which is uniform in n:*

$$P_p(\mathcal{U}, D_n) \geqslant \exp(-\lambda_2 p^{-\alpha_2}).$$

Proof. Let $R(i,j,k)$ be the event that there are α_2 consecutive sites in $\rho(i,j,k)$. By Lemma 27, the probability that D_n is internally spanned is at least

$$\mathbb{P}_p\left(\{D_1 \text{ is i.s.}\} \cap \bigcap_{i=1}^{s_1} \bigcap_{j=1}^{d_1} \bigcap_{k=1}^{n} R(i,j,k) \right).$$

Now we divide $\rho(i,j,k)$ into disjoint blocks of length α_2 to get

$$\mathbb{P}_p(R(i,j,k)) \geqslant 1 - (1-p^{\alpha_2})^{|\rho(i,j,k)|/\alpha_2} \geqslant 1 - \exp\big(- p^{\alpha_2}|\rho(i,j,k)|/\alpha_2 \big).$$

So by Harris's Lemma,

$$P_p(\mathcal{U}, D_n) \geqslant \mathbb{P}_p(D_1 \text{ is i.s.}) \prod_{k=1}^{n} \Big(1 - \exp\big(- p^{\alpha_2}|\rho(i,j,k)|/\alpha_2 \big) \Big)^{\sum_i d_i}.$$

Now, $\mathbb{P}_p(D_1 \text{ is i.s.}) \geqslant p^{c_2}$, where $c_2 = |D_1|$ is a positive constant, so

$$P_p(\mathcal{U}, D_n) \geqslant p^{c_2} \exp\Big(-c_1 \sum_{k=1}^{n} -\log\big(1 - \exp\big(- p^{\alpha_2}|\rho(i,j,k)|/\alpha_2 \big)\big) \Big),$$

where c_1 is the constant $\sum_i d_i$. Property (ii) of Lemma 24 implies that there are positive constants c_3 and c_4 such that $|\rho(i,j,k)| \geqslant c_3 + c_4 k$. Supressing the dependence on n by taking the sum to infinity, we have

$$P_p(\mathcal{U}, D_n) \geqslant \exp\Big(-c_2 \log(1/p) - c_1 \sum_{k=1}^{\infty} -\log\big(1 - \exp\big(- p^{\alpha_2}(c_3 + c_4 k)/\alpha_2 \big)\big) \Big).$$

Let $f : \mathbb{R}^+ \to \mathbb{R}^+$ be the continuous, positive, decreasing, convex function given by

$$f(z) = -\log(1 - e^{-z}).$$

Then

$$\begin{aligned} P_p(\mathcal{U}, D_n) &\geqslant \exp\Big(- c_2 \log(1/p) - c_1 \sum_{k=1}^{\infty} f\big(p^{\alpha_2}(c_3 + c_4 k)/\alpha_2\big)\Big) \\ &\geqslant \exp\left(-c_2 \log(1/p) - \frac{c_1\alpha_2}{c_4} p^{-\alpha_2} \int_0^\infty f(z)\, dz\right) \\ &= \exp\left(-c_2 \log(1/p) - \frac{c_1 \alpha_2 \pi^2}{6c_4} p^{-\alpha_2}\right), \end{aligned}$$

the last equality following from Proposition 5(i) of [16] (although all we actually need is that the integral is finite). ☐

Proof of Theorem 2. First, note that by a standard coupling argument the probability of infecting the origin by time t is increasing in p. Therefore we may assume that p is not just greater than or equal to $(\lambda_2/\log t)^{1/\alpha_1}$, but actually equal to it.

We would like to look inside a canonical quasi-droplet of diameter about t, show that it contains an internally spanned critical droplet, and then show that the infection reaches the origin in time at most t. Unfortunately, the last claim is not quite true: we can only reach the origin in time of the order of $t(\log t)^3$. The solution, as used by Holroyd in [16], is to restrict our attention to a quasi-droplet of diameter about $t/(\log t)^3$. Even with this log-factor, there still exists an internally spanned critical droplet with high probability, but the log-factor gives us just enough extra room to reach the origin by time t.

Let $\tau = t/c_1(\log t)^3$, where c_1 is a constant to be determined later. Let Λ be a minimal canonical quasi-droplet containing the origin, such that $\mathrm{diam}(\Lambda) \geqslant \tau$. Suppose D is a quasi-droplet contained in Λ and let $D_1, D_2, \ldots$ be the sequence of quasi-droplets given by Lemma 24. Let k be minimal such that D_k contains the origin. It is easily verified that there exists a quasi-droplet Λ' containing Λ with $\mathrm{diam}(\Lambda') \leqslant c_2 \,\mathrm{diam}(\Lambda)$, such that no matter how D is chosen, the resulting quasi-droplet D_k is contained in Λ'.

Let E be the event that Λ contains an internally spanned droplet D with $\phi(D) \geqslant p^{-3\alpha_2}$. Noting that there are at least $c_3\tau^2 p^{3\alpha_2}$ disjoint droplets D with $\phi(D) \geqslant p^{-3\alpha_2}$ (since Λ is canonical), it follows by Lemma 29 that

$$\begin{aligned} \mathbb{P}_p(E^c) &\leqslant \big(1 - \exp(-\lambda_2 p^{-\alpha_2})\big)^{c_3\tau^2 p^{3\alpha_2}} \\ &\leqslant \exp\big(- \exp(-\lambda_2 p^{-\alpha_2}) c_3 \tau^2 p^{3\alpha_2}\big) \\ &= \exp\left(- \frac{c_3 \lambda_2^3 t}{c_1^2 (\log t)^9}\right) \to 0. \end{aligned}$$

Let F be the event that, for every $u \in \mathcal{S}_1$ and every $i \in \mathbb{Z}$, every segment of $\ell_u(i)$ of length $p^{-3\alpha_2}$ that is contained entirely within Λ' has a sub-segment of length α_2 that is contained within Π_p. There are at most $c_4\tau^2$ sites in Λ', so

$$\begin{aligned} \mathbb{P}_p(F^c) &\leqslant c_4 s_1 \tau^2 (1 - p^{\alpha_2})^{p^{-3\alpha_2}/\alpha_2} \\ &\leqslant c_4 s_1 \exp\big(2\log\tau - p^{-2\alpha_2}/\alpha_2\big) \\ &\leqslant c_4 s_1 \exp\big(2(\log t - 3\log\log t - \log c_1) - (\log t)^2/(\lambda_2^2 \alpha_2)\big) \to 0. \end{aligned}$$

Claim 30. *The event $E \cap F$ is enough to guarantee that the origin is infected by time t.*

If Claim 30 holds then we are done, since

$$\mathbb{P}_p(E^c \cup F^c) \leqslant \mathbb{P}_p(E^c) + \mathbb{P}_p(F^c) \to 0.$$

Proof of Claim 30. It is sufficient to prove that, given $E \cap F$, the origin is infected in time at most $c_1\tau(\log\tau)^3$, since

$$c_1\tau(\log\tau)^3 \leqslant c_1\frac{t}{c_1(\log t)^3}(\log t)^3 = t.$$

Our strategy will be to prove that there exists a constant c_5 such that the difference between the time D_k becomes fully infected and the time D_{k+1} becomes fully infected is at most $c_5(\log\tau)^3$, for each k. That will be enough to prove the claim, since $0 \in D_{\tau'}$, where $\tau' = c_6\tau$, and $D_{\tau'}$ is infected in time at most

$$|D| + c_5(\log\tau)^3\tau' \leqslant c_7(\log\tau)^3 + c_5c_6\tau(\log\tau)^3,$$

so we would take $c_1 = 2c_5c_6$.

Suppose then, at some time, D_k is fully infected. Define

$$\hat{\xi}(i,j,k) = \ell_{u_i}(i(D_k,u_i)+j) \cap \mathbb{H}_{u_{i-1}}(i(D_k,u_{i-1})+1) \cap \mathbb{H}_{u_{i+1}}(i(D_k,u_{i+1})+1)$$

for $i = 1,\dots,s_1$ and $j = 1,\dots,d_i$. (For $i=1$ we have $\hat{\xi}(i,j,k) = \xi(i,j,k)$, but this is not the case for $i > 1$.) By F, every segment of $\hat{\xi}(i,j,k)$ of length $p^{-3\alpha_2} = ((\log t)/\lambda_2)^3$ has a sub-segment of length α_2 that is infected. By Lemma 23, when $j = 1$, the presence of any one of these sub-segments of length α_2 is enough to infect the whole of $\hat{\xi}(i,1,k)$. The fact that the maximum gap between two length α_2 infected segements is at most $p^{-3\alpha_2}$ means that the infection takes time at most $p^{-3\alpha_2}$. Inducting on j, once $\hat{\xi}(i,j,k)$ is full, by the same argument, $\hat{\xi}(i,j+1,k)$ will be full after at most $p^{-3\alpha_2}$ further steps. Thus the set

$$\bigcup_{i=1}^{s_1}\bigcup_{j=1}^{d_i}\hat{\xi}(i,j,k) \tag{22}$$

will be infected after at most $dp^{-3\alpha_2}$ steps, where $d = \max_i d_i$.

We know from Lemma 27 that

$$D_{k+1} \subset \left[D_k \cup \bigcup_{i=1}^{s_1}\bigcup_{j=1}^{d_i}\hat{\xi}(i,j,k)\right].$$

The sites that are in D_{k+1} but not in (22) can be partitioned into s_1 parallelograms of the form

$$\begin{aligned}&\big(\mathbb{H}_{u_i}(i_{u_i}(D_{k+1})+1) \setminus \mathbb{H}_{u_i}(i_{u_i}(D_k)+1)\big)\\ &\cap \big(\mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D_{k+1})+1) \setminus \mathbb{H}_{u_{i+1}}(i_{u_{i+1}}(D_k)+1)\big).\end{aligned}$$

Each of these parallelograms has size at most d^2, so the infection takes at most d^2 steps.

We have proved that the time between full infection of D_k and full infection of D_{k+1} is at most

$$dp^{-3\alpha_2} + d^2 = O((\log t)^3) = O((\log\tau)^3),$$

as required. ☐

This completes the proof of Theorem 2. ☐

4.6. **Percolation of large quasi-droplets.** For which finite sets Λ is it interesting to talk about percolation on Λ under $\mathcal{U}$ with a random initial set? For example, with two-neighbour bootstrap percolation, one always takes Λ to be a square or a recangle. It would not be interesting to ask whether a random subset of $\Lambda = \{x \in \mathbb{Z}^2 : \|x\|_1 \leqslant n\}$ percolates Λ under the two-neighbour model, since we cannot reach extreme points like $(n, 0)$ without their being initially infected. Thus, if $p = o(1)$, we would have $P_p(\mathcal{U}, \Lambda) = o(1)$ too. In fact, the property that we want Λ to have in order that we avoid situations such as this is exactly the growth property described in the introduction when we discussed the motivation behind the definition of a droplet. For this reason, our results about $P_p(\mathcal{U}, \Lambda)$ will apply only to sets Λ that are quasi-droplets.

With that in mind, in this section we prove analogues of Theorems 1 and 2 for percolation of large quasi-droplets. In keeping with other results of this form in the percolation literature, the bounds that we give are of the same order of magnitude as the corresponding bounds in Theorems 1 and 2.

Theorem 31. *Let $\mathcal{U}$ be critical and let α_1 and λ_1 be as in Theorem 1. Let $\Lambda_1, \Lambda_2, \dots$ be a sequence of quasi-droplets with diameters $d_1, d_2, \dots$ such that $d_n \to \infty$ and let $p_1, p_2, \dots$ be a sequence of probabilities satisfying*

$$p_n \leqslant \left(\frac{\lambda_1}{\log d_n}\right)^{\frac{1}{\alpha_1}}$$

for each $n \geqslant 1$. Then $P_{p_n}(\mathcal{U}, \Lambda_n) \to 0$ as $n \to \infty$.

Proof. Suppose Λ_n is internally spanned. Let Λ'_n be the minimal droplet for $\mathcal{S}$ containing the quasi-droplet Λ_n. We claim that Λ'_n is strongly covered by $A_0^n := \Pi_p \cap \Lambda_n$. Choose a maximal set $\mathcal{B}$ of disjoint breakthrough blocks in A_0^n and run the strong covering algorithm on Λ'_n with $\mathcal{B}$. Let the final droplets be $E_1^n, \dots, E_l^n$ and let $F := A_0^n \setminus E_1^n, \dots, E_l^n$. The closure of $E_1^n \cup \dots \cup E_l^n \cup F$ contains Λ_n (since it contains A_0^n) but the closure of $E_1^n \cup \dots \cup E_l^n$ is itself, so if F is non-empty it must contain a breakthrough block, contradicting the maximality of $\mathcal{B}$. It follows that $\Lambda_n \subset E_1^n \cup \dots \cup E_l^n$, and since the E_i^n are separated and all intersect Λ_n, l must be 1. But E_1 is a droplet and $\Lambda_n \subset E_1 \subset \Lambda'_n$, so $E_1 = \Lambda'_n$. We have proved that Λ'_n is strongly covered by A_0^n.

By Lemma 17, Λ'_n contains a droplet D of diameter between $\log d_n$ and $3 \log d_n$ such that $D \cap \Pi_p$ contains at least $c_1 \log d_n$ breakthrough blocks, for some c_1. By Lemma 20, since $\operatorname{diam}(\Lambda'_n) \leqslant c_2 \operatorname{diam}(\Lambda_n)$ for some c_2, the probability of this is at most $\exp(-c_3 \log d_n)$. □

Theorem 32. *Let $\mathcal{U}$ be critical and let α_2 and λ_2 be as in Theorem 2. Let $\Lambda_1, \Lambda_2, \dots$ be a sequence of canonical quasi-droplets with diameters $d_1, d_2, \dots$ such that $d_n \to \infty$ and let $p_1, p_2, \dots$ be a sequence of probabilities satisfying*

$$p_n \geqslant \left(\frac{\lambda_2}{\log d_n}\right)^{\frac{1}{\alpha_2}}$$

for each $n \geqslant 1$. Then $P_{p_n}(\mathcal{U}, \Lambda_n) \to 1$ as $n \to \infty$.

Proof. The calculation part of the proof follows the corresponding part of the proof of Theorem 2 quite closely, although this time we do not need to use the rescaling trick (that is, we do not need to introduce a new time variable τ). Let E be the event that Λ_n contains an internally spanned droplet D with $\phi(D) \geqslant C p_n^{-3\alpha_2}$, where $C > 0$ is a constant to be determined later in the proof. Noting that there are at least $c_1 d_n^2 p_n^{3\alpha_2}$ disjoint droplets D

with $\phi(D) \geqslant 2p_n^{-3\alpha_2}$, for some constant $c_1 > 0$, it follows by Lemma 29 that

$$\begin{aligned}\mathbb{P}_{p_n}(E^c) &\leqslant \big(1-\exp(-\lambda_2 p_n^{-\alpha_2})\big)^{c_1 d_n^2 p_n^{3\alpha_2}} \\ &\leqslant \exp\big(-\exp(-\lambda_2 p_n^{-\alpha_2})c_1 d_n^2 p_n^{3\alpha_2}\big) \\ &= \exp\left(-\frac{c_1\lambda_2^3 d_n}{(\log d_n)^3}\right) \to 0.\end{aligned}$$

Let F be the event that, for every $u \in \mathcal{S}_1$ and every $i \in \mathbb{Z}$, every segment of $\ell_u(i)$ of length $p_n^{-3\alpha_2}$ that is contained entirely within Λ_n has a sub-segment of length α_2 that is contained within Π_{p_n}. There are at most $c_2 d_n^2$ sites in Λ_n, for some constant $c_2 > 0$, so

$$\begin{aligned}\mathbb{P}_{p_n}(F^c) &\leqslant c_2 s_1 d_n^2\big(1-p_n^{\alpha_2}\big)^{p_n^{-3\alpha_2}/\alpha_2} \\ &\leqslant c_2 s_1 \exp\big(2\log d_n - p_n^{-2\alpha_2}/\alpha_2\big) \\ &= c_2 s_1 \exp\big(2\log d_n - (\log d_n)^2/(\lambda_2^2\alpha_2)\big) \to 0.\end{aligned}$$

Claim 33. *The event $E \cap F$ is enough to guarantee that $\Pi_p \cap \Lambda_n$ percolates Λ_n.*

If Claim 33 holds, then we are done, since

$$\mathbb{P}_{p_n}(E^c \cup F^c) \leqslant \mathbb{P}_{p_n}(E^c) + \mathbb{P}_{p_n}(F^c) \to 0. \tag{23}$$

The proof of the claim is a little less straightforward than one would hope: we cannot simply continue growing from one quasi-droplet to another using the sequence $(D_k)_{k=1}^\infty$ given by Lemma 24, because we encounter problems when we meet one of the sides of Λ. After this point, we find we cannot avoid dealing with shrinking sides. Thus, the majority of the proof of the claim consists of checking that we can continue growing, side by side, once we have first reached the boundary of Λ.

Before we prove the claim, we make the following definition. Given a droplet $D \subset \Lambda$, for each i for which side i of D meets side i of Λ_n (formally, $i_{u_i}(D) = i_{u_i}(\Lambda_n)$), we say side i of D is *active*. For all other values of i, we say side i of D is *dead*.

Proof of Claim 33. We shall prove inductively the existence of a sequence of droplets $D = D_1, D_2, \dots, D_N = \Lambda_n$ such that for $1 \leqslant k \leqslant N-1$, if

$$D_k = \bigcap_{i=1}^{s_1} \mathbb{H}_{u_i}(l_i), \tag{24}$$

then there exists $h \in \{1, \dots, s_1\}$ such that

$$D_{k+1} = \bigcap_{\substack{i=1\\ i\neq h}}^{s_1} \mathbb{H}_{u_i}(l_i) \cap \mathbb{H}_{u_h}(l_h+1). \tag{25}$$

Furthermore, each of these droplets will have the following two properties.

(i) Every side of D_k has Euclidean length at least $2p_n^{-3\alpha_2}$;

(ii) For each $j \geqslant 1$, if exactly j sides of D_k are dead, then each of these sides has Euclidean length at least m_j, where the m_j (which will depend on n) will be chosen later.

We think of D_{k+1} as being obtained by growing D_k along one side. Property (i) ensures that the side of D_k along which we choose to grow contains at least $K + p_n^{-3\alpha_2}$ sites, and that every side of D_k contains at least K sites. Thus by Lemma 26 and event E, if D_k is internally spanned, then D_{k+1} is internally spanned. Since $D_1 = D$ is internally spanned and $D_N = \Lambda_n$, the existence of these droplets will be enough to prove the claim.

Let $\delta > 0$ be a constant with the following property. Whenever we are given quasi-droplets D_k and D_{k+1} as in (24) and (25) such that all the sides of D_k have Euclidean length at least $2p_n^{-3\alpha_2}$, we have

$$|\operatorname{side}_h(D_{k+1})| \geqslant (1-\delta)|\operatorname{side}_h(D_k)|.$$

Suppose we have D_k satisfying (i)-(ii) and that D_k has exactly j dead sides. If $j = 0$ then

$$p_E(D_k) \geqslant p_E(D_{k-1}) \geqslant \ldots \geqslant p_E(D_1) \geqslant s_1 C' p_n^{-3\alpha_2},$$

for some $C' > 0$ (depending on C), where the inequalities follow by noting that the perimeter of D_{k+1} is obtained from the perimeter of D_k by replacing one side by three line segments. So some side h of D_k has Euclidean length at least $C' p_n^{-3\alpha_2}$. The new quasi-droplet D_{k+1} will be that given by (25) with this h. We must check that property (i) holds for D_{k+1} (property (ii) is automatic since $j = 0$). The only side of D_{k+1} which may have shorter length than the corresponding side in D_k is side h. But $|\operatorname{side}_h(D_{k+1})| \geqslant (1-\delta)|\operatorname{side}_h(D_k)| \geqslant (1-\delta)C' p_n^{-3\alpha_2}$, which is certainly larger than $2p_n^{-3\alpha_2}$ for sufficiently large C'.

Now suppose $j \geqslant 1$. We shall prove that there exists an active side h of D_k such that

$$|\operatorname{side}_h(D_k)| \geqslant \frac{\epsilon}{s_1} m_j. \tag{26}$$

for some positive constant ϵ.

We divide the proof into two cases according to whether or not D_k has two non-consecutive dead sides. Before we do that, we need the following observation. Let $I = \{i_1, i_1 + 1 \ldots, i_2 - 1, i_2\}$ be an interval of indices (considered as a subset of $\mathbb{Z}/s_1\mathbb{Z}$) such that $\theta(u_{i_2}) - \theta(u_{i_1}) \leqslant \pi$. Then there exists a unit vector $w \in \mathbb{R}^2$ such that $\langle \operatorname{side}_i(\Lambda_n), w\rangle \geqslant 0$ for all $i \in I$. For fixed I, let $w(I)$ be a unit vector w with this property. There are a finite number of intervals I and for each one there exists $\epsilon(I) > 0$ such that

$$\langle \operatorname{side}_i(\Lambda_n), w(I)\rangle \geqslant \epsilon(I)|\operatorname{side}_i(\Lambda_n)|$$

for all $i \in I$ for which the inner product is strictly positive, so there exists an absolute constant $\epsilon > 0$ such that

$$\langle \operatorname{side}_i(\Lambda_n), w(I)\rangle \geqslant \epsilon|\operatorname{side}_i(\Lambda_n)|$$

for all same range of values of i.

Case 1. Suppose there are two non-consecutive dead sides, so there exist i_1 and i_2, with $i_2 \geqslant i_1 + 2$, such that sides i_1 and i_2 are dead, sides $i_1 + 1, \ldots, i_2 - 1$ are active, and

$$\theta(u_{i_2}) - \theta(u_{i_1}) \leqslant \pi. \tag{27}$$

Let $w = w(\{i_1, \ldots, i_2\})$, so $\langle \operatorname{side}_i(\Lambda_n), w\rangle$ is non-negative for $i_1 \leqslant i \leqslant i_2$ and at least $\epsilon|\operatorname{side}_i(\Lambda_n)|$ for $i_1 + 1 \leqslant i \leqslant i_2 - 1$, and also $\langle \operatorname{side}_i(D_k), w\rangle$ is non-negative for $i_1 \leqslant i \leqslant i_2$ and at least $\epsilon|\operatorname{side}_i(D_k)|$ for $i_1 + 1 \leqslant i \leqslant i_2 - 1$. For $i_1 + 1 \leqslant i \leqslant i_2 - 1$, the u_i sides of D_k and Λ_n coincide, so we have

$$\sum_{i=i_1}^{i_2} \operatorname{side}_i(D_k) = \operatorname{side}_{i_1}(D_k) + \sum_{i=i_1+1}^{i_2-1} \operatorname{side}_i(\Lambda_n) + \operatorname{side}_{i_2}(D_k).$$

By Cauchy-Schwarz,

$$\begin{aligned}\Big|\sum_{i=i_1}^{i_2}\mathrm{side}_i(D_i)\Big| &\geqslant \Big\langle \mathrm{side}_{i_1}(D_k)+\sum_{i=i_1+1}^{i_2-1}\mathrm{side}_i(\Lambda_n)+\mathrm{side}_{i_2}(D_k), w\Big\rangle\\ &= \langle \mathrm{side}_{i_1}(D_k), w\rangle + \sum_{i=i_1+1}^{i_2-1}\langle \mathrm{side}_i(\Lambda_n), w\rangle + \langle \mathrm{side}_{i_2}(D_k), w\rangle\\ &\geqslant \epsilon\sum_{i=i_1+1}^{i_2-1}|\,\mathrm{side}_i(\Lambda_n)|\\ &\geqslant \epsilon(i_2-i_1-1)m_j,\end{aligned}$$

and since $i_2 \geqslant i_1+2$, this last quantity is at least ϵm_j. Therefore by the triangle inequality, there exists $h \in \{i_1+1,\dots,i_2-1\}$ such that

$$|\,\mathrm{side}_h(D_k)| \geqslant \frac{\epsilon m_j}{i_2-i_1-1} \geqslant \frac{\epsilon m_j}{s_1-j} \geqslant \frac{\epsilon}{s_1}m_j,$$

as claimed.

Case 2. Now suppose there is no such pair i_1, i_2. In this case, all the dead sides are consecutive; without loss of generality, they are sides $1,\dots,j$. The failure of (27) means that we must have $\theta(u_1)-\theta(u_j) > \pi$. Let $w = w(\{1,\dots,j\})$, so $\langle \mathrm{side}_i(\Lambda_n), w\rangle$ is non-negative for $1 \leqslant i \leqslant j$ and at least $\epsilon|\,\mathrm{side}_i(\Lambda_n)|$ for $2 \leqslant i \leqslant j-1$, and also $\langle \mathrm{side}_i(D_k), w\rangle$ is non-negative for $1 \leqslant i \leqslant j$ and at least $\epsilon|\,\mathrm{side}_i(D_k)|$ for $2 \leqslant i \leqslant j-1$. This time we have

$$\sum_{i=j+1}^{s_1}\mathrm{side}_i(D_k) = -\Big(\mathrm{side}_1(D_k)+\sum_{i=2}^{j-1}\mathrm{side}_i(\Lambda_n)+\mathrm{side}_j(D_k)\Big),$$

so by Cauchy-Schwarz,

$$\begin{aligned}\Big|\sum_{i=j+1}^{s_1}\mathrm{side}_i(D_k)\Big| &\geqslant \Big\langle \mathrm{side}_1(D_k)+\sum_{i=2}^{j-1}\mathrm{side}_i(\Lambda_n)+\mathrm{side}_j(D_k), w\Big\rangle\\ &= \langle \mathrm{side}_1(D_k), w\rangle + \sum_{i=2}^{j-1}\langle \mathrm{side}_i(\Lambda_n), w\rangle + \langle \mathrm{side}_j(D_k), w\rangle.\end{aligned}$$

Now, since sides 1 and j are dead, we have

$$\min\{|\,\mathrm{side}_1(D_k)|, |\,\mathrm{side}_j(D_k)|\} \geqslant m_j,$$

so

$$\Big|\sum_{i=j+1}^{s_1}\mathrm{side}_i(D_k)\Big| \geqslant j\epsilon m_j \geqslant \epsilon m_j.$$

Hence, by the triangle inequality, there exists $h \in \{j+1,\dots,s_1\}$ such that

$$|\,\mathrm{side}_h(D_k)| \geqslant \frac{\epsilon m_j}{s_1-j} \geqslant \frac{\epsilon}{s_1}m_j,$$

as required.

In both cases we have an active side h of D_k such that (26) holds. The new quasi-droplet will be the D_{k+1} given by (25) with this value of h. We must check properties (i) and (ii) hold for D_{k+1}.

For property (i) we note that, as when $j = 0$, the only side of D_{k+1} that may be shorter than the corresponding side of D_k is the h side, so we require

$$\frac{(1-\delta)\epsilon}{s_1} m_j \geqslant 2p_n^{-3\alpha_2}. \tag{28}$$

For property (ii), dead sides never decrease in length, so we only have to consider what happens when D_{k+1} has one more dead side than D_k. In this case we require

$$\frac{(1-\delta)\epsilon}{s_1} m_j \geqslant m_{j+1} \tag{29}$$

for $j \geqslant 1$, and at the creation of the first dead side we require

$$(1-\delta)C' p_n^{-3\alpha_2} \geqslant m_1. \tag{30}$$

(28) and (29) are simultaneously satisfied by choosing

$$m_j = 2\left(\frac{s_1}{(1-\delta)\epsilon}\right)^{s_1 - j} p_n^{-3\alpha_2}.$$

Finally we can choose C (and hence C') sufficiently large so that (30) is also satisfied. We have now proved the claim. □

This completes the proof of Theorem 32. □

5. Open problems and conjectures

For many update families, the bounds given in Theorems 1 and 2 are far apart because $\alpha_1 \neq \alpha_2$. Which of the two bounds is closer to the truth? One reason why it is hard to answer this question is because there is no precedent: because of the level of generality at which we are working, these are the first results of their kind in which the scale of the threshold isn't necessarily determined exactly. A second reason is that although the strong covering algorithm used to prove Theorem 1 seems to be very generous, and therefore it is tempting to suggest that this is somewhere where the argument could be tightened, the algorithm bears similarities to an algorithm used by Duminil-Copin and Holroyd in [7], in which they determine not only the right scale of the threshold, but also that the threshold is sharp (that is, they prove that one can take $\alpha_1 = \alpha_2$ and $\lambda_1 = \lambda_2$). A third reason is a recent result of Duminil-Copin and van Enter [8] on a so-called anisotropic threshold dynamics model. In this bootstrap process, the neighbourhood $\mathcal{N}$ is

$$\mathcal{N} = \{(1,0),(2,0),(0,1),(-1,0),(-2,0),(0,-1)\}$$

and $\theta = 3$. Theorems 1 and 2 give lower and upper bounds for the critical probability for this threshold process, with $\alpha_1 = 1$ and $\alpha_2 = 2$ respectively. However, Duminil-Copin and van Enter prove that the threshold for infecting the origin by time t is of the order of $p = (\log\log t)^2/\log t$, which is far from both of our bounds.

For many update families $\mathcal{U}$ we do have $\alpha_1 = \alpha_2$. In these cases we conjecture the following.

Conjecture 34. *Suppose that Theorems 1 and 2 hold with constants $\alpha_1 = \alpha_2$. Then there exists a constant $\lambda = \lambda(\mathcal{U})$ such that*

(i) if $\liminf_{t\to\infty} p^{\alpha_1}\log t \geqslant \lambda$, then $P_p(\mathcal{U},t) \to 1$ as $t \to \infty$;

(ii) if $\limsup_{t\to\infty} p^{\alpha_1}\log t \leqslant \lambda$, then $P_p(\mathcal{U},t) \to 0$ as $t \to \infty$.

It is natural to ask whether there are similar results for generalized bootstrap processes in higher dimensions. Given how much more difficult it is to prove sharp threshold results for r-neighbour bootstrap percolation in d dimensions than it is to prove such results for two-neighbour bootstrap percolation in two dimensions, generalizing the results here to higher dimensions may prove quite difficult. Furthermore, we know that in higher dimensions many bootstrap models exhibit a sharp threshold at orders of magnitude very different to powers of $\log t$; for example, r-neighbour bootstrap percolation in $\mathbb{Z}^d$ exhibits a sharp threshold at a power of the $(r-1)$-times iterated logarithm of t (see [4, 5, 2]).

Another interesting question concerns the behaviour of subcritical families. The proof of Theorem 1 goes through for these families with very few modifications (for example, we could just add rules to subcritical families to make new, critical families). This is, of course, not surprising. However, it is interesting to ask what happens from above. For which subcritical families is it true that there exists a function $f(t)$, which is $o(1)$, such that if $p \geqslant f(t)$, then $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$? For which families is it even true that the critical probability for percolation on $\mathbb{Z}^2$ is 0?

In fact, these two questions are equivalent. To see this, first note that the critical probability for percolation on $\mathbb{Z}^2$ is 0 if and only if $\mathbb{P}_p(0 \in [\Pi_p]) = 1$ for all $p > 0$. Now suppose there exists a function $p(t) = o(1)$ such that $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$. In particular, this means that for any fixed $p > 0$ we have $P_p(\mathcal{U}, t) \to 1$ as $t \to \infty$. So, for any $t \geqslant 0$,

$$\mathbb{P}_p(0 \in [\Pi_p]) \geqslant P_p(\mathcal{U}, t) \to 1.$$

Now suppose there exists $p_0 > 0$ such that

$$\mathbb{P}_{p_0}(0 \in [\Pi_{p_0}]) \leqslant 1 - \epsilon,$$

and let $p(t) = o(1)$. For large enough t, we have $p(t) < p_0$, and therefore

$$\limsup_{p \to \infty} \ \mathbb{P}_{p(t)}(0 \in [\Pi_{p(t)}]) \leqslant 1 - \epsilon,$$

meaning that $P_p(\mathcal{U}, t)$ does not tend to 1.

We conclude the section with an open-ended question. It seems that bootstrap percolation on the two-dimensional grid $\mathbb{Z}^2$ may not be the most natural setting for generalized update families. In the arguments above, we rarely used any properties of the grid. In fact, at some points, such as in the proofs of Lemmas 24 and 25, we had to work to overcome the natural structure imposed by $\mathbb{Z}^2$. Is there a more suitable setting in which to study generalized bootstrap processes, for example vertex transitive graphs, or Borel sets in $\mathbb{R}^2$ (as studied by Gravner and Griffeath [11])?

References

1. M. Aizenman and J.L. Lebowitz, *Metastability effects in bootstrap percolation*, J. Phys. A **21** (1988), no. 19, 3801–3813.
2. J. Balogh, B. Bollobás, H. Duminil-Copin, and R. Morris, *The sharp threshold for bootstrap percolation in all dimensions*, Trans. Amer. Math. Soc. **364** (2012), no. 5, 2667–2701.
3. J. Balogh, B. Bollobás, and R. Morris, *Bootstrap percolation in three dimensions*, Ann. Probab. **37** (2009), no. 4, 1329–1380.
4. R. Cerf and E.N.M. Cirillo, *Finite size scaling in three-dimensional bootstrap percolation*, Ann. Probab. **27** (1999), no. 4, 1837–1850.
5. R. Cerf and F. Manzo, *The threshold regime of finite volume bootstrap percolation*, Stochastic Process. Appl. **101** (2002), no. 1, 69–82.
6. J. Chalupa, P.L. Leath, and G.R. Reich, *Bootstrap percolation on a Bethe lattice*, J. Phys. C **12** (1979), no. 1, L31–L35.
7. H. Duminil-Copin and A.E. Holroyd, *Finite volume bootstrap percolation with threshold dynamics on* $\mathbb{Z}^2$ *I: balanced case*, Preprint.

8. H. Duminil-Copin and A.C.D. van Enter, *Sharp metastability threshold for an anisotropic bootstrap percolation model*, Ann. Probab., To appear.
9. C. M. Fortuin, P. W. Kasteleyn, and J. Ginibre, *Correlation inequalities on some partially ordered sets*, Comm. Math. Phys. **22** (1971), 89–103.
10. K. Froböse, *Finite-size effects in a cellular automaton for diffusion*, J. Stat. Phys. **55** (1989), no. 5, 1285–1292.
11. J. Gravner and D. Griffeath, *Threshold growth dynamics*, Trans. Amer. Math. Soc. **340** (1993), no. 2, 837–870.
12. ______, *First passage time for threshold growth dynamics on* $\mathbb{Z}^2$, Ann. Probab. **24** (1996), no. 4, 1752–1778.
13. ______, *Scaling laws for a class of critical cellular automaton growth rules*, Proceedings of the Erdős Center Workshop on Random Walks, 1999, pp. 167–188.
14. J. Gravner, A.E. Holroyd, and R. Morris, *A sharper threshold for bootstrap percolation in two dimensions*, Probab. Theory Related Fields (2010), 1–23.
15. T.E. Harris, *A lower bound for the critical probability in a certain percolation process*, Math. Proc. Cambridge Philos. Soc. **26** (1960), 13–20.
16. A.E. Holroyd, *Sharp metastability threshold for two-dimensional bootstrap percolation*, Probab. Theory Related Fields **125** (2003), no. 2, 195–224.
17. R.H. Schonmann, *On the behaviour of some cellular automata related to bootstrap percolation*, Ann. Probab. **20** (1992), no. 1, 174–193.
18. S. Ulam, *Random processes and transformations*, Proc. Internat. Congr. Math. (1950), 264–275.
19. A.C.D. van Enter, *Proof of Straley's argument for bootstrap percolation*, J. Stat. Phys. **48** (1987), 943–945.
20. J. von Neumann, *Theory of self-reproducing automata*, Univ. Illinois Press, Champaign, 1966.

Trinity College, Cambridge, CB2 1TQ, UK, and Department of Mathematical Sciences, University of Memphis, Memphis, TN 38152, USA
E-mail address: `b.bollobas@dpmms.cam.ac.uk`

Department of Pure Mathematics and Mathematical Statistics, Wilberforce Road, Cambridge, CB3 0WA, UK
E-mail address: `p.j.smith@dpmms.cam.ac.uk`

Department of Mathematical Sciences, University of Memphis, Memphis, TN 38152, USA
E-mail address: `ajuzzell@memphis.edu`